

Surveys for $z > 3$ Damped $\text{Ly}\alpha$ Absorption Systems: the Evolution of Neutral Gas¹

Lisa J. Storrie-Lombardi

SIRTF Science Center, Caltech, MS 100-22, Pasadena, CA 91125, lisa@ipac.caltech.edu

and

Arthur M. Wolfe

*Department of Physics, and Center for Astrophysics and Space Sciences
University of California–San Diego, La Jolla, CA 92093-0424*

Accepted for publication in the *Astrophysical Journal*

ABSTRACT

We have completed spectroscopic observations using LRIS on the Keck 1 telescope of 30 very high redshift quasars, 11 selected for the presence of damped $\text{Ly}\alpha$ absorption systems and 19 with redshifts $z > 3.5$ not previously surveyed for absorption systems. We have surveyed an additional 10 QSOs with the Lick 120" and the Anglo-Australian Telescope. We have combined these with previous data resulting in a statistical sample of 646 QSOs and 85 damped $\text{Ly}\alpha$ absorbers with column densities $N_{\text{HI}} \geq 2 \times 10^{20}$ atoms cm^{-2} covering the redshift range $0.008 \leq z \leq 4.694$. Four main features of how the neutral gas in the Universe evolves with redshift are evident from these data.

1. For the first time we determine a *statistically significant* steepening in the column density distribution function at redshifts $z > 4.0$ ($> 99.7\%$ confidence). The steepening of the distribution function is due to both fewer very high column density absorbers ($N_{\text{HI}} \geq 10^{21}$ atoms cm^{-2}) and more lower column density systems ($N_{\text{HI}} = 2 - 4 \times 10^{20}$ atoms cm^{-2}).
2. The frequency of very high column density absorbers ($N_{\text{HI}} \geq 10^{21}$ atoms cm^{-2}) reaches a peak in the redshift range $1.5 < z < 4$ when the Universe is 10-30% of its present age. Though the sample size is still small the peak epoch appears to be $3.0 \leq z \leq 3.5$. The highest column density absorbers disappear rapidly towards higher redshifts in the range $z = 3.5 \rightarrow 4.7$ and lower redshifts $z = 3.0 \rightarrow 0$. None with column densities $\log N_{\text{HI}} \geq 21$ have yet been detected at $z > 4$ though we have increased the redshift path surveyed $\approx 60\%$.

¹ Some of the observations presented here were obtained at the W.M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation. Observations were also made using the 3.9-m Anglo-Australian Telescope at the Anglo-Australian Observatory and the Shane 3-m Telescope at the Lick Observatory.

3. With our current data set, the comoving mass density of neutral gas, Ω_g , appears to peak at $3.0 < z < 3.5$, but the uncertainties are still too large to determine the precise shape of the Ω_g . The statistics are consistent with a constant value of Ω_g for $2 < z < 4$. There is still tentative evidence for a dropoff at $z > 4$ as indicated by earlier data sets. If we define $R_{g*} \equiv \Omega_g/\Omega_*$, where R_{g*} is the ratio of the peak value of Ω_g to Ω_* , the mass density in galaxies in the local universe, we find values of $R_{g*} = 0.25 - 0.5$ at $z \sim 3$ depending on the cosmology. For an $\Omega = 1$ Universe with a zero cosmological constant $R_{g*} \approx 0.5$. For an $\Omega = 1$ Universe with a positive cosmological constant ($\Omega_\Lambda = 0.7$, $\Omega_M = 0.3$) we find $R_{g*} \approx 0.25$. For a universe with $\Omega_\Lambda = 0$ and $\Omega_M = 0.3$, the value of $R_{g*} \approx 0.3$.
4. Ω_g decreases with redshift for the interval $z = 3.5 \rightarrow 0.008$ for our data set, but we briefly discuss new results from Rao and Turnshek for $z < 1.5$ that suggest $\Omega_g(z < 1.5)$ may be higher than previously determined.

To make the data in our statistical sample more readily available for comparison with scenarios from various cosmological models, we provide tables that includes all 646 QSOs from our new survey and previously published surveys. They list the minimum and maximum redshift defining the redshift path along each line of sight, the QSO emission redshift, and when an absorber is detected, the absorption redshift and measured HI column density.

Subject headings: cosmology—galaxies: evolution—galaxies: quasars—absorption lines

1. Introduction

Surveys for damped Ly α absorption systems have been driven historically by the desire to detect galaxies in very early stages of evolution, before most of the gas has turned in to stars. Because the gas should be neutral and at high column densities ($N_{\text{HI}} > 10^{20}$ atoms cm $^{-2}$) it will leave spectroscopic imprints on the light emitted by background QSOs. These absorbers are identified as damped Ly α absorption systems. Previous studies of the evolution of the neutral gas (Wolfe *et al.* 1986; Lanzetta *et al.* 1991; Wolfe *et al.* 1995 [WLFC95]; Storrie-Lombardi, McMahon & Irwin 1996 [SMI96]), their metal abundances (Lu *et al.* 1996; Pettini *et al.* 1997; Prochaska & Wolfe 1999), their dust content (Fall & Pei 1993; Pei & Fall 1995; Pei, Fall & Hauser 1999), and their kinematics (Prochaska & Wolfe 1997, 1998a, 1998b) provide compelling evidence that damped absorbers are the progenitors of present day galaxies. However, substantial debate continues over exactly which galaxies these are (e.g. Le Brun *et al.* 1997; Haehnelt, Steinmetz & Rauch 1998; Pettini *et al.* 1999; Prochaska & Wolfe 1998a, 1998b; Rao & Turnshek 1998; Salucci & Persic 1999). We present results of a new survey for damped Ly α absorbers in the redshift range $z \approx 3 - 4.5$.

The comoving mass density of neutral gas, $\Omega_g(z)$, at $z \approx 3$ has been known for some time to be comparable to the density of visible matter, i.e. stars in present day galaxies for an $\Omega = 1$ ($\Lambda = 0$) universe (Rao & Briggs 1993; WLFC95; SMI96). We have improved our estimates of $\Omega_g(z)$ at high redshift with new surveys (this work; Storrie-Lombardi & Hook 2000) and estimates of Ω_* have also been updated (Fukugita, Hogan & Peebles 1998). The first damped Ly α survey to have a substantial data set with redshifts $z > 3.0$ (Storrie-Lombardi *et al.* 1996 [SMI96]) hinted at a turnover in $\Omega_g(z)$ at redshifts $z > 4$ (SMI96), prior to which damped Ly α alpha systems might still be collapsing. In this paper we update the discussion of these issues with a larger high redshift data set.

All of the follow-up observations of known candidate damped Ly α absorbers were carried out with the Low Resolution Imaging Spectrometer (LRIS; Oke *et al.* 1995) on the Keck 1 telescope. The signal-to-noise ratios of the resulting spectra are $\approx 25:1$ per pixel and the resolution is $\approx 2 \text{ \AA}$ thus assuring the spectra are adequate for testing the damping hy-

pothesis (cf. WLFC95). The survey data for new damped Ly α candidates was carried out with LRIS on Keck, the KAST spectrograph at the Lick 120", and the RGO spectrograph at the Anglo-Australian Telescope.

The paper is organized as follows. In §2 we present the spectra along with fits of Voigt profiles to the damped Ly α lines to accurately determine the column densities. §3 contains a compilation of a statistical sample of damped Ly α systems including results from all available previous surveys. In §4 we present analyses of the 3 statistical quantities characterizing the evolution of the damped systems, namely, the number of systems per unit redshift interval, dn/dz , the comoving mass density of neutral gas in units of the current critical density, $\Omega_g(z)$, and the frequency distribution of HI column densities, $f(N, z)$. We discuss our results in §5.

2. Data

In this section we first present follow-up spectra of previously reported damped Ly α candidates and then discuss new candidates discovered in our survey.

2.1. LRIS Spectroscopy

Table 1 is a journal of observations describing the LRIS observations of 30 QSOs. Column 1 gives the QSO coordinate name, column 2 the emission redshift, column 3 the UT date of the observations, columns 4 and 5 the V and/or R magnitude of the QSO, column 6 the central wavelength of the grating setting, column 7 the exposure time, column 8 the number of grooves per mm of the grating, and column 9 a reference to observations reporting discovery of the candidate. The spectra were wavelength calibrated with a ThAr lamp, flat fielded with a quartz lamp, and reduced in the usual manner.

2.1.1. Objects Selected for the Presence of Damped Ly α Candidates

Table 2 presents results for the subset of 11 QSOs selected for the presence of damped Ly α candidates. We include only those candidates found in surveys specifically designed to search for strong absorption features blueward of Ly α emission. The confirmed damped Ly α systems are thus included in our statistical sample discussed in § 4 that covers lines of sight observed with sufficient accuracy that damped absorption lines arising in gas with N_{HI} above the

threshold, 2×10^{20} atoms cm^{-2} , could have been detected at the 5σ confidence level. Column 1 gives the QSO coordinate name, column 2 the emission redshift, columns 3 and 4 the V or R magnitude of the QSO, columns 5–8 give the reference to the low resolution survey that identified the candidate absorber, the absorption redshift, the $\text{Ly}\alpha$ equivalent width, and the inferred $\log N(\text{HI})$. Columns 9 and 10 give the absorption redshift and $\log N(\text{HI})$ deduced by profile fitting from the higher resolution LRIS spectra.

Figure 1 exhibits damped $\text{Ly}\alpha$ absorption spectra for the 10 confirmed damped $\text{Ly}\alpha$ systems with $\log N_{\text{HI}} \geq 20.3 \text{ cm}^{-2}$ and the $z = 2.8228$ system toward Q0249–2212 in which $\log N_{\text{HI}} = 20.20 \text{ cm}^{-2}$. Three Voigt profile fits are shown in order of increasing N_{HI} with the middle value representing the mean N_{HI} inferred from the optimal fit, and the other two judged to be $\pm 1 \text{ sigma}$ from the mean. The results of the fits are given in table 2.

Because the high density $\text{Ly}\alpha$ forest in this redshift range can lead to ambiguities in the fitting procedure we did a number of simulations creating spectra with damped absorbers in the forest at known redshifts and column densities, and testing our fitting techniques on these prior to making the final fits on the object spectra. These are extensions of the simulations discussed in SMIH96 where the original damped candidates were selected. The continua of the spectra were first fit with an automated task using cubic splines. These were then modified interactively in dense forest regions where the continuum was placed too low, i.e. the fit had dipped into the line. In every case the redshift of the fit is fixed by the low-ion metal lines detected redward of $\text{Ly}\alpha$ emission; i.e. , outside the $\text{Ly}\alpha$ forest, and a chi square minimization technique, **vpgti**¹ is subsequently used to make the fit.

The metal lines were taken from the LRIS spectra when we had wavelength coverage that went far enough into the red to include them. In the other cases we used the metal line redshifts determined from $5\text{\AA}$ resolution spectra (SMIH96) taken with the ISIS spectrograph on the William Herschel Telescope (WHT). We confirmed that the wavelength calibrations for the WHT and LRIS data agree well in QSOs with overlap regions. The maximum difference in the redshift we determined from the same line measured in different spectra is 0.002 so we are confident that

the metal lines in the WHT spectra don't introduce any significant error when applying the measured redshift to the LRIS spectra.

As expected, the higher resolution of the LRIS spectra make the fitting less ambiguous than the candidate selection process, though the agreement between the estimated and actual column density is very good in almost all the cases. It was crucial to have a low ionization metal line to anchor the redshift for the HI absorption, and in all cases one or both wings of the damped feature provided a good constraint on the column density measurement. Weaker HI components were also used to help constrain the fits, but only the final damped components are shown in the figures.

The following are brief comments on the individual fits:

(1) BR B0019–1522, $z_{\text{abs}} = 3.4370$, $\log N_{\text{HI}} = 20.92$

Si II 1527, Fe II 1608, Al II 1670 were used to determine the redshift. This is the only damped system where the column density estimated from the $5\text{\AA}$ resolution WHT spectrum was substantially in error. This is due to the fact that it lies at $5400\text{\AA}$ which is precisely where the blue and red arms of the ISIS spectra were pasted together.

(2) BRI B0111–2819, $z_{\text{abs}} = 3.1043$, $\log N_{\text{HI}} = 21.0$

Si II 1808 was used to determine the redshift. We did not use the Fe II 1608 and Al II 1670 transitions because they were very broad. We also fitted a weaker component with $z = 3.070$ and $\log N_{\text{HI}} = 19.5 \text{ cm}^{-2}$.

(3) Q0249–2212, $z_{\text{abs}} = 2.8228$, $\log N_{\text{HI}} = 20.2$

Al II 1670 was used to fix the redshift. The fit is excellent owing to absence of $\text{Ly}\alpha$ forest confusion noise near the center of the line profile. Note that the fitted column density, $\log N_{\text{HI}} = 20.2 \text{ cm}^{-2}$ removes this object from the statistical sample.

(4) BR B0951–0450, $z_{\text{abs}} = 3.8580$, $\log N_{\text{HI}} = 20.6$

In common with many other damped $\text{Ly}\alpha$ systems observed at $z > 3$, substantial $\text{Ly}\alpha$ forest structure interferes with the fitting procedure in the wings of the damped feature. In this case we used Si II 1527 observed at low resolution ($\text{FWHM} = 5 \text{\AA}$) with the WHT to anchor the redshift. The column density is constrained by the bump at $5875\text{\AA}$ and the wings on both sides.

(5) BR B0951–0450, $z_{\text{abs}} = 4.2028$, $\log N_{\text{HI}} = 20.4$

We again used Si II 1527 from the WHT spectrum to determine the redshift. The fit is very good due to

¹Written by R.F. Carswell, J.K. Webb, A.J. Cooke, and M.J. Irwin.

the lack of Ly α forest confusion near the line center.

(6) BRI B0952–0115, $z_{abs} = 4.0238$, $\log N_{\text{HI}}=20.55$

A strong C II 1334 line from the LRIS spectrum anchors the fit. The figure shows that the fit is somewhat uncertain owing to strong Ly α absorption in the blue wing of the damped profile. However, the resulting $\log N_{\text{HI}}$ is well determined because the red wing and the blue positive feature near 6085 Å place strong constraints on the fit.

(7) BRI B1013+0035, $z_{abs} = 3.1031$, $\log N_{\text{HI}}=21.1$

We used metal lines from the WHT spectrum to fix the redshift because our limited coverage with LRIS placed all the strong low-ion metal transitions in the Ly α forest. The Ly α forest confusion is especially strong in the blue wing of the damping profile. Nevertheless the fit is again constrained by the observed red wing of the profile and the positive spikes blueward of the profile.

(8) BRI B1114–0922, $z_{abs} = 4.2576$, $\log N_{\text{HI}}=20.3$

There is no LRIS coverage redward of Ly α emission, so we again used metal lines from the WHT spectrum. However, the fit was well constrained by statistically significant positive spikes on both red and blue wings of the damping profile.

(9) BRI B1346–0322, $z_{abs} = 3.7343$, $\log N_{\text{HI}}=20.72$

This damping profile is the best determined of this sample because the redshift is fixed by a single strong C II line in the LRIS spectrum. In addition the absence of significant Ly α forest absorption in the profile wings results in an excellent fit.

(10) BRI B1500+0824, $z_{abs} = 2.7968$, $\log N_{\text{HI}}=20.8$

Because of limited integration time, this is the noisiest spectrum we obtained. However, the rise of the spectrum to the red of the damped Ly α trough places an upper limit not far above the quoted value for N_{HI} . The greater uncertainties are reflected in the larger error bars of the fit. We used WHT metal lines to anchor the redshift, because of our limited LRIS coverage.

(11) BR B2237–0607, $z_{abs} = 4.0691$, $\log N_{\text{HI}}=20.5$

In this case we used C II 1334 which had 2 strong absorption components. Both were used to form the fit shown in figure 1. It is well constrained owing to the high signal-to-noise ratio of the data. This object has since been observed with HIRES (Lu *et al.* 1996) and their result agrees well with ours ($\log N_{\text{HI}} = 20.5$).

2.1.2. New LRIS Survey for Damped Ly α Systems

We obtained LRIS spectra for 19 high redshift QSOs that had not been previously studied for the presence of damped Ly α lines and one QSO with a known damped absorber (Q0201+1120). The objects not previously surveyed come from several quasar samples (Warren, Hewett & Osmer 1991; Schneider, Schmidt & Gunn 1987, 1989 & 1991; McMahon *et al.* 1994; Kennefick *et al.* 1995). Because these objects were not selected on the basis of absorption properties, their spectra are included in constructing the selection function, $g(z)$, which describes the redshift sensitivity of the survey. In cases where damped Ly α lines are detected the spectra are used for fitting Voigt profiles as in the previous section. We show the spectra for all 20 QSOs in figure 2, since even intermediate quality data is lacking for most of these objects. The 1σ error arrays are overplotted, though these are difficult to see in most cases because the signal-to-noise ratio is ≥ 25 . In figure 3 we show the spectra for the confirmed damped systems. The damped candidate and confirmed absorbers are discussed individually below.

(1) Q0046–293, $z_{abs} = 3.0589$, $\log N_{\text{HI}}=20.1$

The redshift was determined from narrow Si II 1526, Fe II 1608, and Al II 1670 and the fit is well constrained by the steep sides of the HI trough. This system is not included in the statistical sample as it falls below the threshold of $\log N_{\text{HI}}=20.3 \text{ cm}^{-2}$.

(2) Q0046–293, $z_{abs} = 3.841$, $\log N_{\text{HI}}=20.1$

This system is composed of two components, with the redshifts determined from C II 1334. It is also well constrained by the HI line. This system is not included in the statistical sample as it falls below the threshold of $\log N_{\text{HI}}=20.3 \text{ cm}^{-2}$.

(3) Q0057–274, $z_{abs} = 3.2413$, $\log N_{\text{HI}}=19.8$

The redshift was determined from C II 1334 and the fit is well constrained by the sides of the HI trough. This system is not included in the statistical sample as it falls below the threshold of $\log N_{\text{HI}}=20.3 \text{ cm}^{-2}$.

(4) PSS J0059+0003, $z_{abs} = 3.1043$, $\log N_{\text{HI}}=20.0$

The redshift is well determined from Si II 1526. Though there is strong Ly α absorption in the blue wing of the damped profile the resulting $\log N_{\text{HI}}$ is accurate because the red wing places a strong constraint on the fit.

(5) Q0201+1120, $z_{abs} = 3.3848$, $\log N_{\text{HI}}=21.3$

The damped system in this QSO was studied by White, Kinney & Becker (1993) in a spectrum with $\approx 10\text{\AA}$ resolution. We have wavelength coverage from 5000–8000 $\text{\AA}$ which includes several low ionization metal lines from the damped absorber. Our column density and redshift measurements are in excellent agreement with those previously published.

- (6) PC0953+4749, $z_{abs}=3.403$, $\log N_{\text{HI}} \leq 20.9$
 $z_{abs}=3.890$, $\log N_{\text{HI}} \leq 21.1$

We originally observed this object because of the obvious large absorption features visible in the published spectrum. We did not plan to include it in the statistical sample because (a) the QSO was approximately a magnitude fainter than the objects we were originally following up and (b) we observed it because we knew it might have high column damped features. When we completed our new survey, we had observed and included in our statistical sample all of the QSOs with $V \leq 19.5$ and $z \geq 3.5$ that were available at the time. This object has a published V magnitude of 19.5, so we now include it in our statistical sample. It is problematic as it is one of the faintest objects we observed and we only have wavelength coverage at one blue setting. The existing QSO spectra have 25 $\text{\AA}$ resolution which is too low to provide any useful metal line information. Even though the 1.5 $\text{\AA}$ resolution LRIS spectrum has sufficient signal-to-noise to fit the HI we are unable to accurately anchor the redshifts for these absorbers so the column densities quoted are upper limits.

- (7) PSS J1435+3057, $z_{abs}=3.26$, $\log N_{\text{HI}}=20.0$
 $z_{abs}=3.51$, $\log N_{\text{HI}}=20.0$
 $z_{abs}=3.71$, $\log N_{\text{HI}}=20.0$

Previous experience shows that these candidates are highly unlikely to be damped. They are not included in the statistical sample in the analysis that follows.

- (8) PSS J1443+2724, $z_{abs}=4.216$, $\log N_{\text{HI}}=20.8$

Though taken with the 300 g/mm grating, the signal-to-noise ratio in this spectrum is high enough to determine that this absorber is highly likely to be damped. The redshift was determined from the strong, narrow C II 1334 line.

- (9) PC2047+0123, $z_{abs} = 2.7299$, $\log N_{\text{HI}}=20.4$

Although the only useful low-ion metal lines, C II 1334 and Fe II 1608, were in the Ly α forest portion of the LRIS spectrum, we believe the redshift is well determined, since neither metal line was severely blended.

(10) Q2050+359, NOT DAMPED

Though the feature at $\approx 4843\text{\AA}$ appears to be damped at first glance, an examination of the associated metal lines reveals that it is not. There is a complex of low and high ionization metals lines that cover the redshift range $2.98 \leq z \leq 3.17$, and there is no suitable fit for a damped Ly α feature with $\log N_{\text{HI}} \geq 20.0$. It is a complex of lower column density lines.

2.1.3. New Lick/AAT Survey for Damped Ly α Systems

Ten additional QSOs with redshifts $z \geq 4$ were observed at the Lick 3-m telescope with the KAST spectrograph or the Anglo-Australian Telescope with the RGO spectrograph, at $\approx 6\text{\AA}$ resolution. The survey data from these objects are summarized in table 4 and the spectra are shown in figure 4. PSS J0244–0108 is excluded from further analysis because it exhibits broad absorption lines believed to be intrinsic to the quasar. Four damped candidates were detected in these spectra and their estimated column densities were determined from the measured equivalent widths of the lines. This method is discussed in detail in Lanzetta *et al.* 1991 and its application to $z > 4$ QSOs is discussed in SMIH96. BRI B0111–2819 was subsequently followed up with LRIS and is included in table 2.

3. Statistical Sample

The statistical sample of damped absorbers includes all confirmed or candidate damped Ly α absorbers with neutral hydrogen column densities $N_{\text{HI}} \geq 2 \times 10^{20}$ atoms cm^{-2} from surveys designed to detect these absorbers (Wolfe *et al.* 1986; Lanzetta *et al.* 1991; LWT95; WLFC95; SMIH96; Storrie-Lombardi, Irwin & McMahon [SIM96]; Storrie-Lombardi & Hook 2000; this paper). The QSOs with damped absorbers are listed in table 5. The QSO spectra that did not contain damped absorbers, and the redshift path they contribute, are listed in table 6. It includes 646 QSO lines of sight and 85 damped Ly α absorbers. The spectra for all of these lines of sight had sufficient signal-to-noise to detect damped absorption with N_{HI} above the threshold at the 5σ confidence level between the redshifts z_{min} and z_{max} listed in the tables. The redshift path surveyed in each QSO is used to construct the sensitivity function, $g(z)$. It gives the number of lines of sight at a given redshift over which

damped systems can be detected at a $> 5\sigma$ level. The form of the selection function is given by

$$g(z) = \sum_{i=1}^m \Theta(z_i^{max} - z) \Theta(z - z_i^{min}) \quad (1)$$

where Θ is the Heaviside step function and where the sum is over the m QSOs in the survey (see Lanzetta *et al.* 1991).

The upper panel of figure 5 illustrates how the redshift sensitivity for the entire statistical sample compares with previous work. The data included in this paper are shown as a solid line, the large compilation of damped systems in the Large Bright Quasar Survey (WLFC95) which provides the bulk of the data for $z < 3$ is shown as a dashed line, and the APM survey (SMIH96) which provided most of the previous data for $z > 3$ is shown as a dotted line. The lower panel of figure 5 shows the detail for the redshift range $z \geq 3$.

The new data included in this paper increase by 75 percent the redshift path surveyed for $z \geq 3$. In table 7 we compare Δz for the LBQS survey, the APM survey, and this paper for $z > 3$ and for the entire redshift range of the samples. We also give analogous values for the total “absorption distance” ΔX which is defined as

$$X(z) = \begin{cases} \frac{2}{3}[(1+z)^{3/2} - 1] & \text{if } q_0 = 0.5; \\ \frac{1}{2}[(1+z)^2 - 1] & \text{if } q_0 = 0. \end{cases} \quad (2)$$

(Bahcall & Peebles 1969; cf. Tytler 1987). The redshift path over which damped systems could be detected is crucial in estimating the cosmological mass density in neutral gas from the damped systems, as well as the number density per unit redshift, and the frequency distribution per unit column density. These are discussed in detail in the next section.

4. Analysis

4.1. Frequency Distribution of Column Densities, $f(N, z)$

By combining our statistical sample of damped systems with the data presented in Hu *et al.* (1995) for column densities $N_{\text{HI}} \leq 2 \times 10^{20}$ atoms cm^{-2} , we can examine the frequency distribution for the entire HI column density range, $10^{12} - 10^{22}$ atoms cm^{-2} (see Tytler 1987; Petitjean *et al.* 1993; Hu *et al.* 1995). This is shown in figure 6. To first order it is fit by a power law function, $f(N) \propto N^{-1.46}$, over ten orders of magnitude in column density. The Ly α forest

absorbers are by far the most numerous, yet if we integrate this to find the mass density, the bulk of the HI mass is locked up in the damped systems. A disadvantage of a power law model for the HI column density distribution of damped Ly α absorbers is the divergent nature of the integral mass contained in the systems. A high column density cutoff (e.g. 10^{22}) must be selected to do the integral. We discuss below an alternative parameterization based on a gamma distribution (c.f. Pei & Fall 1995; SIM96).

4.1.1. Differential Column Density Distribution

The frequency distribution of HI column densities is defined as follows. Let $f(N, X)dNdX$ be the number of absorbers per sightline with HI column densities in the interval $(N, N + dN)$, and absorption distances in the interval $(X, X + dX)$. Then the frequency distribution of HI column densities $f(N, z) \equiv f(N, X[z])$. In figure 7 we show the $f(N, z)$ derived from the entire statistical sample of damped absorbers for $q_0 = 0.0$, and in figure 8 we show the same data divided into four redshift bins: $z = 0.008 - 1.5$, $z = 1.5 - 2.5$, $z = 2.5 - 3.5$, and $z = 3.5 - 4.7$. The flattest distribution is in the redshift range $z = 3.0 - 3.5$, and it steepens towards higher and lower redshifts from this point. This was noted in WLFC95 for the $z < 3.5$ data and in SIM96 for the $z > 3.5$ data, but the differences in $f(N)$ with redshift were not statistically significant. Our new statistical sample is large enough to make some quantitative statements about the redshift evolution of $f(N)$ at high redshift.

4.1.2. The Neutral Gas Density at $z < 1.5$

There is a steepening of the differential distribution of column densities evident at $z < 1.5$ in our data set but not at a statistically significant level. Previous surveys have shown a very lower number density per unit redshift from International Ultraviolet Explorer and Hubble Space Telescope (HST) data (LWT95) and Jannuzi *et al.* (1998), also find numbers consistent with this from HST Key Project data. But Rao & Turnshek (1998) have reported in a Letter the discovery of 12 new low redshift damped systems and their full survey has just appeared in preprint form (Rao & Turnshek 2000). This redshift range is problematic because any bias in QSO surveys due to possible obscuration by dust in foreground absorbers would have the largest impact here (see Pei & Fall 1995; SMI96). We focus in this paper on the redshift range $z > 1.5$.

4.1.3. Evolution of the Neutral Gas Density at $z > 1.5$

For the first time we determine a *statistically significant* ($> 99.7\%$ confidence) steepening in the column density distribution function at redshifts $z > 4.0$. This is evidenced by comparing the data in the redshift range $1.5 < z < 4$ with the data at $z \geq 4$ as shown in figure 9. This shows the cumulative distribution, normalized by the absorption distance surveyed. A Kolmogorov-Smirnov (K-S) test gives a probability of only 0.006 that the two redshift samples are drawn from the same distribution. The steepening of the distribution function is due to both fewer very high column density absorbers ($N_{\text{HI}} \geq 10^{21}$ atoms cm^{-2}) and more lower column density systems ($N_{\text{HI}} = 2 - 4 \times 10^{20}$). No damped systems with column densities $\log N_{\text{HI}} \geq 21$ have yet been detected at $z > 4$.

4.1.4. Fits to the Column Density Distribution

A single power law, $f(N) = kN^{-\beta}$ does not provide a good fit to the column density distribution function for damped Ly α absorbers. A single power law fit has the additional problem that if $\beta < 2$, as all current estimates indicate, then the total mass in damped systems diverges unless an upper bound to the HI column density is assumed. An alternative parameterization using a gamma function to describe the HI column density distribution was suggested by Pei & Fall (1995) and adopted by SIM96. We use the same formalism here.

We model the data with a gamma distribution of the form

$$f(N, z) = (f_*/N_*)(N/N_*)^{-\beta} e^{-N/N_*} \quad (3)$$

where f_* is the characteristic number of absorbing systems at the column density N_* , and N_* is a parameter defining the turnover, or ‘knee’, in the number distribution. Both f_* and N_* may in general vary with redshift but for the moment we treat them as constants. This functional form is similar to the Schechter luminosity function (Schechter 1976). For $N \ll N_*$ the gamma function tends to the same form as the single power law, $f(N) \propto N^{-\beta}$; while for $N \gtrsim N_*$, the exponential term begins to dominate. We use a maximum likelihood technique to find a solution over a two-dimensional grid of pairs of values of N_* and β , since the constant f_* can be directly

computed using the constraint

$$m = \sum_{i=1}^n f_* \int_{N_{\min}}^{N_{\max}} \int_{z_{\min}^i}^{z_{\max}^i} f(N, z) dz dN \quad (4)$$

where m is the total number of observed systems.

The results of fits to the data in the redshift ranges $1.5 < z < 4$ and $z \geq 4$, overplotted on the cumulative distribution of absorbers, are shown in the upper panels of figure 10. A data point for the expected number of Lyman-limit systems that would be detected down to $\log N_{\text{HI}} = 17.2$ is shown with a circled star. Lyman-limit systems are defined and detected by the observation of neutral hydrogen (HI) absorption which is optically thick to Lyman continuum radiation for $\lambda < 912\text{\AA}$, the Lyman limit, corresponding to a column density $N(\text{HI}) \geq 1.6 \times 10^{17} \text{ cm}^{-2}$ (See Tytler 1982; Sargent, Steidel & Boksenberg 1989; Lanzetta 1991; Storrie-Lombardi *et al.* 1994; Stengler-Larrea *et al.* 1995 for discussions of Lyman Limit systems.) Their contribution is calculated by integrating the number density per unit redshift of Lyman limit systems expected ($N(z) = 0.27(1+z)^{1.55}$) over the redshift path covered by the n QSOs in the damped sample, i.e.

$$\begin{aligned} \text{LLS}_{\text{expected}} &= \sum_{i=1}^n \int N(z) dz = \sum_{i=1}^n \int_{z_{\min}}^{z_{\text{em}}} N_0(1+z)^{\gamma} dz \\ &= \sum_{i=1}^n \int_{z_{\min}}^{z_{\text{em}}} 0.27(1+z)^{1.55} dz \end{aligned} \quad (5)$$

(SMIH94, SIM96). Including the Lyman limit system point provides a longer baseline in column density and allows us to make better estimates of the uncertainties in the fit. The log-likelihood function results with confidence contours are shown in the lower panels of figure 10.

The parameters for the fit in equation 3 are very similar for both redshift ranges when the Lyman limit systems are included, with $\log N_* = 21.22$ atoms cm^{-2} for both and $\beta = 1.24$, $f_* = 0.05$ for the lower redshift data and $\beta = 1.08$, $f_* = 0.16$ for the higher redshift data. Without the Lyman-limit point anchoring the lower column density end, the fits to just the high column density damped systems yield $\beta = 1.52 \pm 0.25$, $f_* = 0.03$, and $\log N_* = 21.39 \pm 0.30$ for the lower redshift data and $\beta = 0.32 \pm 0.6$, $f_* = 0.34$, and $\log N_* = 20.92 \pm 0.50$ for the higher redshift data. The errors on the exponent β are very large for the z

> 4 data due to the small number of data points. The fits reflect what we saw in looking at the differential counts, i.e. the “knee”, N_* , where the distribution turns over is at a lower column density for the higher redshift data. The best fit results are tabulated in table 8.

In SIM96, with a much smaller data set (366 QSOs, 44 absorbers), we obtained $\log N_* = 21.63 \pm 0.35$, $\beta = 1.48 \pm 0.30$, and $f_* = 1.77 \times 10^{-2}$ fitting the whole sample. With our current statistical sample (646 QSOs, 85 absorbers) we find very similar results with $N_* = 21.33 \pm 0.30$, $\beta = 1.48 \pm 0.25$, and $f_* = 3.19 \times 10^{-2}$. Alternatively, single power law fits of the form $f(N) \propto N^{-\beta}$ (not including the point for the Lyman-limit systems) yield $\beta = 1.96$ for $z > 4$ and $\beta = 1.78$ for $1.5 < z < 4$, again consistent with values obtained in earlier work (LWTLMH91; SIM96).

4.2. Number Density Evolution with Redshift

We now examine the redshift evolution of the damped Ly α absorbers in our combined data set by determining the number density of absorbers per unit redshift, $dN/dz \equiv N(z)$. In a standard Friedmann Universe for absorbers with cross-sectional area A , absorption distance X , and Φ_0 per unit comoving volume,

$$N(z) = \Phi_0 A c H_0 dX/dz \quad (6)$$

where

$$dX/dz = \left[\frac{(1+z)^2}{(1+z)^2(1+\Omega_M z) - z(z+2)\Omega_\Lambda} \right]^{1/2}. \quad (7)$$

It is customary to represent the number density as a power law of the form

$$N(z) = N_0(1+z)^\gamma, \quad (8)$$

The differential distribution in number density of absorbers versus redshift is shown in figure 11. It is fit by a single power law with $N_0 = 0.055$ and $\gamma = 1.11$, which would suggest no intrinsic evolution in the product of the space density and cross-section of the absorbers with redshift, but the fit is very poorly constrained. This is evident in the $>68.3\%$ and $>95.5\%$ confidence contours for the log-likelihood results when calculating γ and N_0 which is shown in figure 12. The poor fit is due to differential evolution in the number density of damped Ly α absorbers with different column densities which has been described in LWT95, WLFC95, and SIM96.

The number density of absorbers versus redshift, split at a column density of $\log N_{\text{HI}} = 21$ is shown in figure 13. The number density of systems with $\log N_{\text{HI}} > 21$, shown as solid lines, peaks at $z \approx 3.5$, when the Universe is 15-20% of its present age. These systems then disappear at a much faster rate from $z=3.5$ to $z=0$ than does the population of damped absorbers as a whole. There is a paucity of very high column density systems at the highest redshifts surveys, which was also evidenced in the steepening of the column density distribution discussed in § 4.1.3. The number density of damped absorbers with column densities $\log N_{\text{HI}} \leq 21$ decreases from redshifts $z \approx 4$ to $z \approx 3.5$ and remains relatively constant towards $z = 1.5$. The differential evolution with column density suggests:

1. There has been insufficient time at $z > 3.5$ for the highest column density absorbers to collapse.
2. Once they do form, the highest column density absorbers preferentially form stars before their lower column density counterparts, and hence disappear more rapidly towards lower redshifts.

From the evolution of the HI with redshift alone we are unable to determine if we are watching the evolution of similar systems with redshift or watching some systems disappear and others form.

4.3. Evolution of the Mass Density of Neutral Gas

The comoving mass density of neutral gas is given by

$$\Omega_g(z) = \frac{H_0}{c} \frac{\mu m_H}{\rho_{\text{crit}}} \frac{\sum_i N_i(\text{HI})}{\Delta X(z)} \quad (9)$$

where the density is in units of current critical density (see Lanzetta *et al.* 1991). The quantity μ is the mean particle mass per m_H where the latter is the mass of the H atom, ρ_{crit} is the current critical mass density, and the sum is over damped Ly α systems in the statistical sample listed in table 5. We use $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}$ in all these calculations. Because we are interested in determining $\Omega_g(z)$ for separate redshift bins, $\Delta X(z)$ is given by the integral $\int g(X) dX$ between the $X(z)$ corresponding to the redshift limits of each bin. We determine $\Omega_g(z)$ by assuming $\mu = 1.3$. The results for our data set from $z = 0.008$ to $z = 4.7$ are shown in figure 14 for 3 different cosmologies with $\Omega = \Omega_M + \Omega_\Lambda$:

- $\Omega_M=1, \Omega_\Lambda=0$
- $\Omega_M=0.3, \Omega_\Lambda=0$
- $\Omega_M=0.3, \Omega_\Lambda=0.7$

The region Ω_* is the $\pm 1\sigma$ range for the mass density in stars in nearby galaxies (Fukugita, Hogan & Peebles 1998). The point at $z=0$ is the value inferred from 21 cm emission from local galaxies (Zwaan *et al.* 1997). These results are tabulated in table 9. Ω_g evolves in the same fashion with redshift regardless of the cosmology selected and there is a tentative peak in Ω_g in the redshift range $3 < z < 3.5$. We consistently find very high column density systems in this redshift range, even though for $z > 4$ QSOs, $z \sim 3$ absorbers are deep in the Ly α forest where they are most difficult to detect due to the density of the forest lines in the spectra. This was evidenced in section § 4.1.1 where the shape of the frequency distribution of column densities was flattest in the $3 < z < 3.5$ redshift bin.

What does change with cosmological model is the ratio of the peak value of Ω_g to Ω_* , the mass density in galaxies in the local universe. We define this ratio as $R_{g*} \equiv \Omega_g(3 < z < 3.5)/\Omega_*$. Previous surveys (Lanzetta *et al.* 1991; WLFC95; SMI96) have found the value of R_{g*} to be of order 1. When we look at this ratio ($R_{g*} = \Omega_g(z=3)/\Omega_*$) For our current data set and value of Ω_* , we find $R_{g*} \approx 0.25 - 0.5$, depending on the cosmology. For an $\Omega = 1$ Universe with a zero cosmological constant $R_{g*} \approx 0.5$, for an $\Omega = 1$ Universe with a positive cosmological constant ($\Omega_\Lambda = 0.7, \Omega_M = 0.3$) $R_{g*} \approx 0.25$, and for a universe with $\Omega_\Lambda = 0$ and $\Omega_M = 0.3$, $R_{g*} \approx 0.3$.

5. Discussion

Though the results for the comoving mass density of neutral gas shown in figure 14 look very similar to those published in previous work we discuss below the main differences in these new results and their implications.

1. The values of Ω_g in this work are lower than those previously published for $z > 1.5$. The values for Ω_g published in WLFC95 and SMI96 used $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}$ and in this work we use $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}$ which decreases Ω_g by 23% at all redshifts. The improved statistics due to the increased redshift path also reduces the effect of the scatter produced by the small

number of QSOs with very high column density absorbers. The value of Ω_g is now less sensitive to the maximum column density in any particular redshift bin. The comparison of Ω_g to Ω_* does not appear very different, even though the values are lower, because our estimate of Ω_* has changed as well.

2. The value of Ω_* from Fukugita, Peebles & Hogan (1998) that we are using in these plots is also about half the value inferred from Gnedin & Ostriker (1992) used in previous determinations. There are still substantial uncertainties in the relative contributions of disks and spheroids in determining the local mass density (e.g. Salucci & Persic 1999; J. Dalcanton, R. Marzke and L. Yan, private communication). Salucci & Persic (1999) find a lower value of Ω_* (2×10^{-3}) independent of the Hubble constant, which is about half the value we have plotted from Fukugita, Peebles & Hogan (1998). As our understanding of the local mass density improves we will continue to refine these numbers for comparison with the high redshift determinations.
3. The calculation of Ω_g for $z > 3.5$ in SMI96 hinted at a turnover at these high redshifts. Our larger data set does not confirm whether or not this turnover is real though the trend continues to be present in our larger dataset. We do see a statistically significant change in the frequency distribution of column densities at $z > 4$ (§ 4.1.3), which suggests we are probing an epoch when the highest column density absorbers have not yet collapsed. Also, for one of the highest column density high redshift systems ($\log N_{\text{HI}} \leq 21.1$ at $z = 3.89$) from PC0953+4749, we only have an upper limit for the column density as discussed in section (§ 2.1.1). Though we have surveyed enough quasars at $z > 3.5$ to not be extremely sensitive to the inclusion or exclusion of one absorber, it is true that if this high column density were substantially lower, the error bars for the $3 < z < 3.5$ and $z > 3.5$ bins would not overlap. We will have to wait for more data to make a definitive statement about a turnover in Ω_g at $z > 4$.
4. Ω_g determined in WLFC95 shows a monotonic decrease from $z = 3.5 \rightarrow 0.008$. Our larger data set also shows a decrease, but the inclusion of

the addition survey of Storrie-Lombardi & Hook (2000) makes Ω_g flatter in the redshift $2 < z < 3$ range. Our new data are consistent with a constant value of Ω_g for $2 < z < 4$. New results from Rao and Turnshek (2000) for $z < 1.5$ also suggest $\Omega_g(z < 1.5)$ may be higher than previously determined. In figure 15 we replot the first panel of figure 14, and show their lower redshift data point with dashed lines. The error bars are still very large but they find a substantially higher value for Ω_g at $z < 1.65$ than previous surveys.

We thank Julia Kennefick, George Djorgovski, Mike Irwin, and Richard McMahon for providing QSO coordinates prior to publication and the staff at Keck, Lick, and the AAT for their able assistance in obtaining these observations. LSL thanks Jason Prochaska, Celine Peroux, and Max Pettini, and an anonymous referee for helpful suggestions.

REFERENCES

- Bahcall J.N., Peebles P.J.E. 1969, *ApJ*, 156, L7
- Fall, S.M. & Pei, Y.C. 1993, *ApJ*, 402, 479
- Fukugita, M., Hogan, C.J., & Peebles, P.J.E. 1998, *ApJ*, 503, 518
- Gnedin, N.Y. & Ostriker, J.P. 1992, *ApJ*, 400, 1
- Haehnelt, M.G., Steinmetz, M. & Rauch, M. 1998, *ApJ*, 495, 647
- Hu E.M., Kim T., Cowie L.L., Songaila A. & Rauch M. 1995, *AJ*, 110, 1526
- Jannuzi, B.T., Bahcall, J.N., Bergeron, J., Bok-
senberg, A., Hartig, G.F. Kirhakos, S., Sargent,
W.L.W., Savage, B.D., Schneider, D.P., Turnshek,
D.A., Weymann, R.J., & Wolfe, A.M. 1998, *ApJS*,
118, 1
- Kennefick, J.D., de Carvalho, R.R., Djorgovski, S.G.,
Wilber, M.M., Dickson, E.S., Weir, N., Fayyad, U.
& Roden, J., 1995, *AJ*, 110, 78
- Lanzetta K.M., 1991, *ApJ*, 375, 1
- Lanzetta K.M., Wolfe A.M., Turnshek D.A., Lu L.,
McMahon R.G., Hazard C., 1991, *ApJS*, 77, 1
- Lanzetta K.M., Wolfe A.M., Turnshek D.A., 1995,
ApJ, 440, 435 [LWT95]
- Le Brun, V., Bergeron, J., Boisse, P. & Deharveng,
J.M. 1997, *A&A*, 321, 733
- Lu, L., Sargent, W.L.W., Womble, D.S. & Barlow,
T.A. 1996, *ApJ*, 457, L1
- McMahon, R.G., Omont, A., Bergeron, J., Kreysa,
E., & Haslam, C.G.T., 1994, *MNRAS*, 267, L9
- Natarajan, P. & Pettini, M. 1997, *MNRAS*, 291, 28
- Oke, J.B., Cohen, J.G., Carr, M., Cromer, J., Din-
gizian, A., Harris, F.H., Labrecque, S., Lucinio,
R., Schall, W., Epps H., and Miller, J. 1995, *PASP*,
107, 375
- Pei, Y.C. & Fall, S.M. 1995, *ApJ*, 454, 69
- Pei, Y.C., Fall, S.M. & Hauser, M.G. 1999, *ApJ*, 552,
604
- Petitjean P., Webb J.K., Rauch M., Carswell R.F. &
Lanzetta K.M. 1993 *MNRAS*, 262, 499
- Pettini M., Smith L.J., Hunstead R.W., King D.L.,
1994, *ApJ*, 426, 79
- Pettini M., Smith L.J., King D.L., Hunstead R.W.
1997, *ApJ*, 486, 665
- Pettini M., Ellison, S.L., Steidel, C.C. & Bowen, D.V.
1999, *ApJ*, 510, 576
- Prochaska, J.X. & Wolfe, A.M. 1997, *ApJ*, 487, 73
- Prochaska, J.X. & Wolfe, A.M. 1998a, *ApJ*, 494, 15
- Prochaska, J.X. & Wolfe, A.M. 1998b, *ApJ*, 507, 113
- Prochaska, J.X. & Wolfe, A.M. 1999, *ApJS*, 121, 369
- Rao, S. & Briggs, F. 1993, *ApJ*, 419, 515
- Rao, S.M. & Turnshek, D.A. 1998, *ApJ*, 500, L115
- Sargent, W.L.W., Steidel, C.C., & Boksenberg, A.
1989, *ApJS*, 79, 703
- Salucci, P. & Persic, M. 1999, *MNRAS*, 309, 923
- Schneider, D.P., Schmidt, M. & Gunn, J.E., 1987,
ApJ, 316, L1
- Schneider, D.P., Schmidt, M. & Gunn, J.E., 1989, *AJ*,
98, 1507

- Schneider, D.P., Schmidt, M. & Gunn, J.E., 1991, *AJ*, 101, 2004
- Stengler-Larrea, E.A., Boksenberg, A., Steidel, C.C., Sargent, W.L.W., Bahcall, J.N., Bergeron, J., Hartig, G.F., Jannuzi, B.T., Kirhakos, S., Savage, B.D., Schneider, D.P., Turnshek, D.A., & Weymann, R.J., 1995, *ApJ*, 444, 64
- Storrie-Lombardi, L.J., McMahon, R.G., Irwin, M.J. & Hazard, C. 1994, *ApJ*, 427, L13
- Storrie-Lombardi, L.J., McMahon, R.G., Irwin, M.J. & Hazard, C. 1996, *ApJ*, 468, 128 [SMIH96]
- Storrie-Lombardi, L.J., McMahon, R.G. & Irwin, M.J. 1996, *MNRAS*, 283, L79 [SMI96]
- Storrie-Lombardi, L.J., Irwin, M.J., and McMahon, R.G. 1996, *MNRAS*, 282, 1330 [SIM96]
- Storrie-Lombardi, L.J. & Hook, I.M. 2000, *ApJS*, submitted
- Tytler D. 1982, *Nature*, 298, 427
- Tytler D. 1987, *ApJ*, 321, 49
- Warren S.J., Hewett P.S. & Osmer P.S., 1991, *ApJS*, 76, 23
- White, R.L., Kinney, A. & Becker, R.H. 1993, *ApJ*, 407, 456
- Wolfe, A.M. 1987, *Proc. Phil. Trans. Roy. Soc.*, 320, 503
- Wolfe, A.M., Lanzetta, K.M., Foltz C.B., & Chaffee F.H., 1995, *ApJ*, 454, 698 [WLFC95]
- Wolfe, A.M., Turnshek, D.A., Smith, H.E., & Cohen, R.D. 1986, *ApJS*, 61, 249
- Zwaan, M.A., Briggs, F.H., Sprayberry, D., & Sorar, E. 1997, *ApJ*, 490, 173


























Fig. 1.— The spectra for the 10 confirmed damped Ly α systems with $\log N_{\text{HI}} \geq 20.3 \text{ cm}^{-2}$ and the $z = 2.8228$ system toward Q0249–2212 in which $\log N_{\text{HI}} = 20.20 \text{ cm}^{-2}$ are plotted. Three Voigt profile fits are shown in order of increasing N_{HI} with the middle value representing the mean N_{HI} inferred from the optimal fit, and the other two judged to be $\pm 1\sigma$ from the mean. The results are summarized in table 2.






Fig. 2.— All of the spectra taken in the new LRIS survey, plus Q0201+1120, are shown here. The 1σ error arrays are overplotted as dashed lines, though these are difficult to see in most cases due to the high signal-to-noise ratios of the spectra. The observations are summarized in table 3. Expanded versions of the region around the confirmed damped systems are shown in figure 3.




Fig. 3.— The confirmed damped Ly α systems with $\log N_{\text{HI}} \geq 20.3 \text{ cm}^{-2}$ from the new survey are plotted. Three Voigt profile fits are shown in order of increasing N_{HI} with the middle value representing the mean N_{HI} inferred from the optimal fit, and the other two judged to be $\pm 1\sigma$ from the mean. The results are summarized in table 3.














Fig. 4.— The survey spectra taken at the Lick 3-m and the Anglo-Australian Telescope are shown. The data are summarized in table 4.



as a dotted line. The lower panel shows the detail for the redshift range $z \geq 3$. The new data included in this paper nearly double the redshift path previously surveyed for $z \geq 3$.

Fig. 5.— The redshift sensitivity function, $g(z)$, is plotted for the combined data sets used in this survey as a solid line. It gives the number of lines of sight at a given redshift over which damped systems can be detected at a $> 5\sigma$ level. The upper panel illustrates how the redshift sensitivity for the entire statistical sample compares with previous work. The data included in this paper are shown as a solid line, the large compilation in the LBQS survey (WLFC95) which provides the bulk of the data for $z < 3$ is shown as a dashed line, and the APM survey (SMIH96) which provided most of the previous data for $z > 3$ is shown




Fig. 6.— The column density distribution function of neutral hydrogen for $12 \leq \log N_{\text{HI}} \leq 22$ the damped Ly α absorbers. To first order it is fit by a power law, $f(N) \propto N^{-1.46}$, over 10 orders of magnitude in column density.




Fig. 7.— The differential column density distribution function of neutral hydrogen determined for the entire statistical sample of damped Ly α absorbers.




Fig. 8.— The differential column density distribution function of neutral hydrogen for damped Ly α absorbers shown in four redshifts bins: 0.008-1.5, 1.5-2.5, 2.5-3.5, and 3.5-4.7. The distribution is flattest at $z \approx 3$ and steepens towards higher and lower redshifts from that point. The steepening at high redshift is due to both a paucity of very high column density systems as well as an increase in the frequency of lower column density systems. There is also a steepening of the differential distribution evident at $z < 1.5$ but this is not statistically significant (see Rao & Turnshek 2000 for new results in this redshift range).




Fig. 9.— The log of the cumulative number of damped Ly α absorbers versus log column density normalized by the redshift path surveyed, is shown versus log column density for two redshift ranges. The solid line shows the data for $1.5 < z < 4.0$ and the dashed line shows the data for $z \geq 4.0$. A Kolmogorov-Smirnov (K-S) test gives a probability of only 0.006 that the two redshift samples are drawn from the same distribution. This is the first time that the steepening of the column density distribution function has been shown to be statistically significant.




Fig. 10.— The left hand plots show $1.5 < z < 4.0$ and the right hand plots show $z \geq 4.0$. In the top two panels we show the log of the cumulative number of damped Ly α absorbers versus log column density, including a point for the expected number of Lyman limit systems as a circled star. The best fits to the column density distribution function of the form $f(N, z) = (f_*/N_*)(N/N_*)^{-\beta}e^{-N/N_*}$ are overplotted using the values in table 8. In the lower two panels we show the corresponding $>68.3\%$, $>95.5\%$, and $>99.7\%$ confidence contours for the log-likelihood results when calculating β , the slope of the power law portion of the fit, and $\log N_*$, the knee in the distribution.




Fig. 11.— The differential distribution in number density of absorbers versus redshift for the entire data set is shown. It is fit by a single power law with $N_0 = 0.055$ and $\gamma = 1.11$, which would suggest no intrinsic evolution in the product of the space density and cross-section of the absorbers with redshift, but the fit is very poorly constrained as shown in figure 12.




Fig. 12.— The $>68.3\%$ and $>95.5\%$ confidence contours are plotted for the log-likelihood results for γ and N_0 in fitting the number density per unit red-

shift with a single power law $N(z) = N_0(1+z)^\gamma$. The fit for a single power law is very poorly constrained due to differential evolution in the number density of damped Ly α absorbers with different column densities. This is illustrated in figure 13.




Fig. 13.— The number density per unit redshift is shown for the same data shown in figure 11, but now split into two groups at a column density of $\log N_{\text{HI}} = 21$. The number density of systems with $\log N_{\text{HI}} > 21$, shown as solid lines, peaks at $z \approx 3.5$, when the Universe is 15-20% of its present age. These systems then disappear at a much faster rate from $z=3.5$ to $z=0$ than does the population of damped absorbers as a whole. There is a paucity of very high column density systems at the highest redshifts surveys, which was also evidenced in the steepening of the column density distribution discussed in § 4.1.3. The number density per unit redshift of damped absorbers with column densities $\log N_{\text{HI}} \leq 21$ peaks at $z \approx 4$, drops at $z \approx 3.5$ and remains constant or increases slightly towards $z = 1.5$.






Fig. 14.— The comoving mass density in neutral gas contributed by damped Ly α absorbers, $\Omega_g(z)$, is plotted for three different cosmologies: (a) $\Omega_M=1$, $\Omega_\Lambda=0$, (b) $\Omega_M=0.3$, $\Omega_\Lambda=0$, and (c) $\Omega_M=0.3$, $\Omega_\Lambda=0.7$. The region Ω_* is the $\pm 1\sigma$ range for the mass density in stars in nearby galaxies (Fukugita, Hogan & Peebles 1998). The point at $z=0$ is the value inferred from 21 cm emission from local galaxies (Zwaan *et al.* 1997). These results are tabulated in table 9.




Fig. 15.— This is the first panel ($\Omega_M=1$, $\Omega_\Lambda=0$) of figure 14, showing the comoving mass density in neutral gas contributed by damped Ly α absorbers, $\Omega_g(z)$, with the value from Rao & Turnshek (2000) for $z < 1.65$ plotted as dashed lines. The error bars are still very large, but they find a substantially higher value of Ω_g at lower redshifts than indicated from previous surveys.

TABLE 1
LRIS JOURNAL OF OBSERVATIONS

QSO	z_{em}	Date	V	R	$\lambda_{central}$ (Å)	Exp (secs)	Grating (g/mm)	Reference
BR BB0019–1522	4.528	18 Sep 1995		19.0	5865 7065 7765	3300 1200 1500	1200 1200 1200	1
Q 0046–293	4.014	19 Sep 1995	19.4		5390 6565	2400 1200	1200 1200	2
Q 0057–274	3.52	20 Sep 1995		18.73	5085 6260	1500 600	1200 1200	2
PSS J0059+0003	4.16	19 Sep 1995		19.5	5190 6365	2400 1200	1200 1200	3
PC 0104+0215	4.171	20 Sep 1995	19.7		5390 6565	3600 2400	1200 1200	4
BRI B0111–2819	4.30	20 Sep 1995 20 Sep 1995 19 Sep 1995 19 Sep 1995		18.7	5390 6865 5190 6365	4300 1200 2400 1200	1200 1200 1200 1200	5
PC 0131+0120	3.792	20 Sep 1995		19.4	5665 6865	1500 1200	1200 1200	6
Q 0201+1120	3.61	20 Sep 1995 19 Sep 1995 19 Sep 1995 19 Sep 1995	20.1		6500 5665 6565 7565	1600 1800 1500 1200	300 1200 1200 1200	7
PSS J0248+1802	4.43	18 Sep 1995		18.4	5190 6365 7565	1700 1200 600	1200 1200 1200	3
Q 0249–222	3.20	18 Sep 1995	18.4		5190 6365	1200 500	1200 1200	8
PC 0345+0130	3.638	20 Sep 1995 19 Sep 1995	19.9		5190 6260	2400 1500	1200 1200	6
PC 0751+5622	4.281	7 Mar 1995	19.7		6150	3600	1200	4
BR B0951–0450	4.369	7 Mar 1995		18.9	6150	2800	1200	1
BRI B0952–0115	4.426	7 Mar 1995		18.7	6150	2200	1200	1
PC 0953+4749	4.46	7 Mar 1995	19.3		5450	5400	1200	9
BRI B1013+0035	4.405	7 Mar 1995		18.8	5450	3200	1200	1
BRI B1114–0822	4.495	7 Mar 1995		19.4	6150	3600	1200	1
PSS J1317+3531	4.365	16 May 1996		19.1	6500	622	300	3
BRI B1346–0322	3.992	7 Mar 1995		19.1	6150	2800	1200	1
PSS J1430+2828	4.306	16 May 1996		19.3	6500	600	300	3
PSS J1435+3057	4.297	16 May 1996		19.3	6500	600	300	3
PSS J1443+2724	4.407	16 May 1996		19.3	6500	600	300	3
MG 1500+0431	3.67	20 Sep 1995		18.01	5085 6260	1200 600	1200 1200	10

TABLE 1—*Continued*

QSO	z_{em}	Date	V	R	$\lambda_{central}$ (Å)	Exp (secs)	Grating (g/mm)	Reference
BRI B1500+0824	3.943	7 Mar 1995		19.3	5100	1800	1200	1
PC 1548+4637	3.544	20 Sep 1995	19.2		5085	2400	1200	9
					6260	600	1200	
PC 1640+4628	3.700	19 Sep 1995	19.5		5085	1800	1200	9
		20 Sep 1995			6260	2100	1200	
PC 2047+0123	3.799	18 Sep 1995	19.7		5085	5400	1200	9
					6260	1800	1200	
Q 2050–359	3.49	19 Sep 1995	18.3		5085	1200	1200	2
					6260	600	1200	
BR B2237–0607	4.558	18 Sep 1995		18.3	6565	2400	1200	1
					7765	1200	1200	
PC 2331+0216	4.093	19 Sep 1995	20.0		5665	2800	1200	4
					6865	1200	1200	

REFERENCES.— (1) Storrie-Lombardi *et al.* 1996a (2) Warren, Hewett & Osmer 1991 (3) Kennefick *et al.* 1995 (4) Schneider, Schmidt & Gunn 1989 (5) McMahon & Irwin, private communication (6) Schneider, Schmidt & Gunn 1987 (7) White, Kinney & Becker 1993 (8) Sargent, Steidel & Boksenberg 1989 (9) Schneider, Schmidt & Gunn 1991 (10) McMahon *et al.* 1994

TABLE 2
LRIS FOLLOW-UP OF CANDIDATE DAMPED LYMAN- α ABSORBERS

QSO	QSO Properties			Ref.	Survey Data			Follow-up Properties	
	z_{em}	V	R		z_{abs} (Å)	W (cm ⁻²)	$\log N_{\text{HI}}$	z_{abs} (cm ⁻²)	$\log N_{\text{HI}}$
BR B0019–1522	4.528		19.0	1	3.42	7.6	20.0	3.4370	20.92±0.10
				1	3.98	12.3	20.5	not damped	
				1	4.28	8.0	20.1	not damped	
BRI B0111–2819	4.30		18.7	2	3.100	23.4	21.0	3.1043	21.0±0.1
Q 0249–2212	3.202	18.4		3	2.831	10.2	...	2.8228	20.2±0.1
BR B0951–0450	4.369		18.9	1	3.84	24.0	21.0	3.8477	20.6±0.1
				1	4.20	10.6	20.3	4.2028	20.4±0.1
BRI B0952–0115	4.426		18.7	1	4.01	18.0	20.8	a 4.0221	20.2
								b 4.0255	20.3
					components ‘a’ and ‘b’ combined →			4.0238	20.55±0.10
BRI B1013+0035	4.405		18.8	1	3.10	17.5	20.8	3.1031	21.1±0.1
				1	3.73	9.6	20.2	not damped	
BRI B1114–0822	4.495		19.4	1	3.91	6.7	19.9	not damped	
				1	4.25	11.7	20.4	4.2576	20.3±0.1
				1	4.45	5.3	19.7	not damped	
BRI B1346–0322	3.992		19.1	1	3.73	10.0	20.3	3.7343	20.72±0.10
BRI B1500+0824	3.943		19.3	1	2.80	11.3	20.4	2.7968	20.8±0.2
BR B2237–0607	4.558		18.3	1	4.08	11.5	20.4	a 4.0683	20.2
								b 4.0699	20.25
					components ‘a’ and ‘b’ combined →			4.0691	20.5±0.1

REFERENCES.— (1) Storrie-Lombardi *et al.* 1996a; (2) this paper (3) Sargent, Steidel & Boksenberg 1989

TABLE 3
NEW LRIS SURVEY DATA

QSO	z_{em}	z_{min}	z_{max}	z_{abs}	W (Å)	$\log N_{\text{HI}}$ (cm^{-2})
Q 0046–293	4.014	2.882	3.964	3.0589		20.1±0.1
				3.8407		(a)19.9±0.1
				3.8422		(b)19.6±0.1
components ‘a’ and ‘b’ combined →				3.841		20.1±0.1
Q 0057–274	3.52	2.603	3.475	3.2413		19.8±0.1
PSS J0059+0003	4.16	2.750	4.108	3.1043		20.0±0.1
PC 0104+0215	4.171	2.881	4.119			
PC 0131+0120	3.792	3.116	3.744			
Q 0201+1120	3.61	...	...	3.3848		21.3±0.1
PSS J0248+1802	4.43	2.810	4.376			
PC 0345+0130	3.638	2.699	3.592			
PC 0751+5623	4.281	3.526	4.228			
PC 0953+4749	4.457	3.010	4.004	3.403		≤20.9
				3.890		≤21.1
PSS J1317+3531	4.365 [†]	2.978	4.311			
PSS J1430+2828	4.306 [†]	2.777	4.253			
PSS J1435+3057	4.297 [†]	2.905	4.244	3.26	7.6	20.0
				3.51	7.5	20.0
				3.71	7.3	20.0
PSS J1443+2724	4.407 [†]	2.950	4.353	4.216	18.1	20.8
MG 1500+0431	3.67	2.606	3.623			
PC 1548+4637	3.544	2.607	3.499			
PC 1640+4628	3.700	2.604	3.653			
PC 2047+0123	3.799	2.620	3.751	2.7299	11.2	20.4±0.1
Q 2050–359	3.49	2.605	3.445	2.98	7.8	not damped
PC 2331+0216	4.093	3.115	4.042			

NOTE.—The confirmed systems have uncertainties quoted for the column density measurements. The candidate systems do not. Previous experience shows that the system in PSS J1443+2724 is highly likely to be damped, and the candidates in PSS J1435+3057 are highly unlikely to be damped. All the candidate and confirmed systems with $\log N_{\text{HI}} \geq 20.3$ are included in the statistical sample in the analysis that follows, except for Q 0201+1120. It was observed because of a known damped absorber, not in an unbiased survey.

[†]The emission redshifts for these QSOs have been determined from the mean redshifts of the OI 1302, SiIV+OIV 1400, and CIV 1549 emission lines observed in the LRIS spectra.

TABLE 4
NEW LICK/AAT SURVEY DATA

QSO	Ref.	R	Where	Date	Exp (Seconds)	z_{em}	z_{min}	z_{max}	z_{abs}	W (Å)	$\log N_{\text{HI}}$ (cm^{-2})
PSS J0030+1702	1	19.3	Lick	24 Oct 95	10800	4.282	2.763	4.229			
BRI B0046−2458	2	18.9	AAT	27 Aug 95	3600	4.15	2.575	4.099			
BRI B0111−2819	2	18.7	AAT	27 Aug 95	3295	4.30	2.908	4.247	3.100	23.4	21.0 ^b
PSS J0117+1552	1	18.6	Lick	24 Oct 95	5400	4.244	2.646	4.192			
PSS J0132+1341	1	19.3	Lick	24 Oct 95	7200	4.147	2.844	4.096	3.93	10.5	20.3
BRI B0135−4239	2	18.5	AAT	27 Aug 95	4200	3.97	2.575	3.920			
PSS J0244−0108 ^a	1	19.0	Lick	24 Oct 95	3300	4.00					
BR B0331−1622	2	17.9	Lick	23 Oct 95	9000	4.40	2.868	4.326	3.56	14.6	20.6
PSS J1057+4555	1	17.7	Lick	20,23 Oct 95	4500	4.101	2.652	4.050	3.05	10.2	20.3
BR B1600+0729	2	18.8	Lick	16 Mar 96	4200	4.38	3.062	4.326			

NOTE.—The Lick observations were taken at the 120" with the KAST spectrograph with simultaneous blue and red arm observations. The Anglo-Australian Telescope observations were taken with the RGO spectrograph. All are at approximately 6Å resolution (FWHM). For the spectra with enough wavelength coverage to the red to see CIV, the QSO emission redshifts have been determined from a mean of the OI 1302, SiIV+OIV 1400, and CIV 1549 emission lines.

^aThis QSO exhibits broad absorption lines and is excluded from further consideration.

^bThis damped candidate was subsequently observed with LRIS (see table 2 and figure 2).

REFERENCES.— (1) Kennefick et al. 1995 and J. Kennefick, private communication (2) M. Irwin, private communication

TABLE 5
STATISTICAL SAMPLE: QSOs WITH DAMPED LYMAN-ALPHA ABSORBERS

QSO	z_{em}	z_{min}	z_{max}	z_{abs}	$\log N_{\text{HI}}$ (cm^{-2})	Refs.
Q 0000–2619	4.11	2.389	4.060	3.3901	21.4	5,13
Q 0010–0012	2.15	1.634	2.119	2.0233	20.8	4,10
Q 0013–0029	2.08	1.634	2.049	1.9730	20.7	4,11
BR B0019–1522	4.528	2.97	4.473	3.4370	20.92	22,1
Q 0027+0103	2.29	1.634	2.257	1.9375	20.6	4,10
Q 0042–2930	2.39	1.591	2.354	1.931	20.5	4
Q 0049–2820	2.26	1.638	2.223	2.0713	20.5	4,11
Q 0056+0125	3.16	2.197	3.119	2.7750	21.0	4,10
Q 0058–2914	3.07	1.778	3.052	2.6711	21.2	21
Q 0100–3105	2.64	1.687	2.605	2.131	20.5	4
Q 0100+1300	2.69	1.64	2.74	2.3093	21.4	16,15
Q 0102–1902	3.04	2.044	2.995	2.3693	21.0	5,8
Q 0102–0214	1.98	1.649	1.949	1.7431	20.6	4,10
BRI B0111–2819	4.30	2.709	4.247	3.1043	21.0	1
Q 0112–3041	2.99	1.881	2.945	2.4191	20.5	21,9
–	–	–	–	2.7023	20.3	21,9
Q 0112+0300	2.81	1.813	2.785	2.4227	21.0	5,11
PSS J0132+1341	4.147	2.844	4.096	3.93	20.3	1
Q 0149+3335	2.43	1.64	2.43	2.1413	20.5	16,7
Q 0201+3634	2.49	1.632	2.879	2.4614	20.4	21,8
–	–	–	–	1.768	20.5	21
Q 0216+0803	3.00	1.731	2.953	2.2930	20.5	5,9
BR B0331–1622	4.38	2.868	4.326	3.56	20.6	1
Q 0336–0142	3.20	2.109	3.155	3.0619	21.2	21,8
Q 0347–3819	3.23	2.044	3.186	3.0244	20.8	21,18
Q 0449–1330	3.097	2.006	3.056	2.052	20.4	21
Q 0458–0203	2.29	1.96	2.29	2.0399	21.7	16,7
Q 0528–2505	2.779	1.961	2.741	2.1404	21.0	5
Q 0834–2006	2.75	1.632	2.704	1.715	20.4	21
Q 0836+1122	2.70	1.74	2.67	2.4660	20.6	16,6
Q 0913+0715	2.78	1.866	2.739	2.6187	20.3	21,8
MG 0930+2858	3.41	2.173	3.366	3.24	20.5	2
Q 0935+4143	1.9800	1.0626	1.550	1.369	20.3	3
BR B0951–0450	4.369	2.93	4.315	3.8580	20.6	22,1
–	–	–	–	4.2028	20.4	22,1
BRI B0952–0115	4.426	2.99	4.372	4.0238	20.55	22,1
PC 0953+4749	4.457	3.010	4.004	3.403	20.9	1
–	–	–	–	3.890	21.1	1
BRI B1013+0035	4.405	2.61	4.351	3.1031	21.1	22,1
Q 1032+0414	3.39	2.067	3.347	2.839	20.3	21
PSS J1057+4555	4.101	2.652	4.050	3.05	20.3	1
BRI B1108–0747	3.922	2.64	3.873	3.607	20.33	22,23
BRI B1114–0822	4.495	3.19	4.440	4.2576	20.3	22,1

TABLE 5—*Continued*

QSO	z_{em}	z_{min}	z_{max}	z_{abs}	$\log N_{\text{HI}}$ (cm^{-2})	Refs.
Q 1151+0651	2.76	1.65	2.76	1.7737	21.3	16,6
Q 1159+0132	3.27	1.988	3.226	2.6846	21.1	21,8
BR B1202–0725	4.694	3.16	4.637	4.383	20.49	22,23
Q 1205+0918	2.08	1.634	2.046	1.673	20.6	4
Q 1209+0919	3.30	2.175	3.254	2.5835	21.4	21,8
Q 1210+1731	2.54	1.634	2.502	1.8920	20.6	4,10
Q 1215+3322	2.61	1.64	2.60	1.9989	21.0	16,7
Q 1223+1753	2.92	1.945	2.879	2.4658	21.5	4,11
Q 1232+0815	2.57	1.789	2.534	2.3376	20.9	4,10
Q 1240+1516	2.28	1.634	2.247	1.738	20.7	4
Q 1244+3443	2.48	1.64	2.50	1.8593	20.5	16,7
Q 1246–0217	2.11	1.634	2.075	1.779	21.2	4
Q 1308+0105	2.80	1.634	2.763	1.762	20.6	4
GB 1320+3927	2.98	1.968	2.940	2.11	20.4	2
Q 1329+4117	1.9350	0.4853	0.6318	0.5193	20.7	3
Q 1337+1121	2.92	1.86	2.92	2.7957	20.9	16,6
BRI B1346–0322	3.992	2.65	3.942	3.7343	20.72	22,1
Q 1347+1116	2.70	1.92	2.71	2.4709	20.3	16,6
Q 1409+0930	2.86	1.979	2.800	2.4561	20.5	21,8
PSS J1443+2724	4.407	2.950	4.353	4.216	20.8	1
Q 1451+1223	3.26	2.158	3.207	2.478	20.4	16,21
BR BI1500+0824	3.943	2.39	3.894	2.7968	20.8	22,1
GB 1610+2806	3.54	2.021	3.498	2.59	20.6	2
MG 1614+0506	3.21	1.984	3.168	2.52	20.4	2
GB 1759+7539	3.05	1.955	3.010	2.624	20.77	2
PC 2047+0123	3.799	2.620	3.751	2.7299	20.4	1
Q 2112+0555	0.4660	0.1105	0.4513	0.2039	20.4	3
Q 2132–4321	2.42	1.595	2.386	1.916	20.7	4
Q 2138–4427	3.17	2.107	3.128	2.851	20.9	4,12
Q 2206–1958	2.56	1.85	2.58	1.9205	20.5	16,14
–	–	–	–	2.0763	20.7	16,17
Q 2223–0512	1.4040	0.4159	0.6310	0.4925	20.9	3
–	–	0.9259	1.3800	–	–	3
Q 2230+0232	2.15	1.634	2.119	1.8642	20.8	4,10,11
Q 2231–0015	3.015	1.749	2.980	2.0657	20.6	4,9
BR B2237–0607	4.558	2.96	4.502	4.0691	20.5	22,1
Q 2239–3836	3.55	2.389	3.508	3.2810	20.8	21,9
Q 2248+0127	2.56	1.634	2.524	1.9080	20.6	4,10
Q 2348–0108	3.01	2.044	2.965	2.4272	20.5	16,6
–	–	–	–	2.6161	21.3	21,6
Q 2351+0217	2.03	1.634	2.000	1.766	20.9	4,10
Q 2359–0216	2.31	1.747	2.779	2.0951	20.7	16,7
–	–	–	–	2.1537	20.3	16,7

REFERENCES.—See table 6 for reference list.

TABLE 6
STATISTICAL SAMPLE: QSOs WITHOUT DAMPED
LYMAN-ALPHA ABSORBERS

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
Q 0001+0842	3.241	2.024	3.229	21
Q 0002+151	1.8990	0.4723	0.6034	3
—	—	1.1198	1.5500	3
Q 0003+158	0.4500	0.0080	0.4355	3
MG 0004+1359	3.25	1.899	3.207	2
Q 0004+1711	2.898	2.002	2.851	21
Q 0006+0230	2.09	1.787	2.059	4
Q 0006+0200	2.35	1.634	2.317	4
Q 0007−0004	2.26	1.634	2.227	4
Q 0007−000	2.29	1.670	2.260	16
MG 0007+0141	2.90	1.882	2.861	2
Q 0007+106	0.0890	0.0080	0.0781	3
Q 0009−0138	1.99	1.634	1.960	4
Q 0009+0219	2.66	1.784	2.623	4
Q 0009−0215	2.11	1.634	2.079	4
Q 0014+8118	3.380	1.928	3.340	21
Q 0014−0256	1.85	1.729	1.821	4
Q 0015+0239	2.47	1.784	2.435	4
Q 0016+0045	2.31	1.651	2.277	4
Q 0018−0220	2.56	1.634	2.524	4
Q 0018+0047	1.83	1.655	1.802	4
Q 0020+0217	1.80	1.665	1.772	4
Q 0022+0150	2.77	1.791	2.732	4
Q 0023+0010	1.90	1.657	1.871	4
Q 0025−0151	2.08	1.634	2.049	4
Q 0026+0158	1.89	1.727	1.861	4
Q 0026+129	0.1420	0.0080	0.1306	3
Q 0027+0149	2.33	1.694	2.297	4
Q 0028+0236	2.00	1.634	1.970	4
Q 0028−0148	2.08	1.840	2.049	4
Q 0029−0152	2.39	2.013	2.356	4
PSS J0030+1702	4.282	2.763	4.229	1
Q 0037−018	2.34	1.654	2.303	16
Q 0039−2630	1.81	1.634	1.782	4
Q 0040−2917	2.09	1.634	2.056	4
Q 0041−2638	3.045	1.657	3.029	21
Q 0041−2707	2.79	1.668	2.748	4
Q 0041−2607	2.79	1.634	2.470	4
Q 0041−2658	2.46	1.634	2.422	4
Q 0041−2859	2.13	1.589	2.103	4
Q 0042−3053	1.97	1.634	1.944	4
Q 0042−2627	3.298	2.113	3.253	21

TABLE 6—*Continued*

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
Q 0042–2656	3.33	2.215	3.314	21
Q 0042–2657	2.90	2.226	2.859	4
Q 0043–2937	2.23	1.656	2.198	4
Q 0044+030	0.6240	0.3386	0.6078	3
Q 0045–3002	2.02	1.603	1.991	4
Q 0045–0341	3.138	1.961	3.094	21
Q 0045–013	2.53	1.784	2.493	16
Q 0046–293	4.014	2.882	3.964	1
BRI B0046–2458	4.15	2.575	4.099	1
Q 0047–2759	2.13	1.649	2.099	4
Q 0047–3050	2.97	1.930	2.933	4
Q 0047–2538	1.97	1.591	1.939	4
Q 0047–2326	3.422	2.291	3.378	21
Q 0048–0119	1.88	1.634	1.849	4
Q 0048–2545	2.08	1.634	2.051	4
Q 0049–0104	2.10	1.715	2.065	4
Q 0049–0012	1.95	1.634	1.916	4
Q 0049+007	2.27	1.644	2.238	16
Q 0049+014	2.31	1.681	2.276	16
Q 0049+171	0.0640	0.0080	0.0534	3
Q 0050+124	0.0611	0.0080	0.0505	3
Q 0050–2523	2.16	1.592	2.127	4
Q 0051–0226	2.53	1.634	2.491	4
Q 0052–0058	2.21	1.634	2.180	4
Q 0052+251	0.1550	0.0080	0.1435	3
Q 0053–0134	2.06	1.634	2.031	4
Q 0053–2824	3.616	2.454	3.576	21
Q 0054+0200	1.87	1.634	1.844	4
Q 0054+144	0.1710	0.0080	0.1593	3
Q 0054–006	2.76	1.854	2.724	16
Q 0055+0141	2.23	1.651	2.200	4
Q 0055–2744	2.20	1.567	2.163	4
Q 0055–2629	3.6560	1.920	3.609	21
Q 0055–0200	1.98	1.782	1.953	4
Q 0055+0025	1.91	1.634	1.885	4
Q 0056–0241	2.23	1.779	2.194	4
Q 0057–0225	2.01	1.715	1.979	4
Q 0057–274	3.52	2.603	3.475	1
Q 0058–2604	2.47	1.606	2.437	4
Q 0058–0227	2.23	1.712	2.194	4
Q 0058+0155	1.95	1.634	1.924	4
Q 0059–0207	2.29	1.653	2.257	4
Q 0059–2625	2.10	1.614	2.069	4
Q 0059+0035	2.55	1.673	2.510	4
PSS J0059+0003	4.16	2.750	4.108	1

TABLE 6—*Continued*

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
Q 0100+0146	1.91	1.692	1.880	4
Q 0101–2548	1.97	1.596	1.943	4
Q 0101–3025	4.073	1.937	3.116	21
Q 0102–0240	1.84	1.731	1.818	4
BRI B0103+0032	4.437	2.87	4.383	22
Q 0103–0141	2.21	1.634	2.174	4
Q 0103–2901	2.87	1.922	2.831	4
Q 0104+0030	1.87	1.667	1.845	4
PC 0104+0215	4.171	2.881	4.119	1
Q 0105–2649	2.46	1.667	2.428	4
Q 0106–0230	2.28	1.634	2.246	4
Q 0106+0119	2.10	1.871	2.068	4
Q 0107+0022	1.97	1.634	1.938	4
Q 0108+0028	2.01	1.733	1.975	4
Q 0109+022	2.35	1.734	2.317	16
Q 0110–0107	1.89	1.643	1.860	4
Q 0112–2728	2.894	1.784	2.855	21
Q 0114–0856	3.163	1.838	3.118	21
Q 0115–3002	3.249	1.733	3.207	21
PSS J0117+1552	4.244	2.646	4.192	1
Q 0117+213	1.4930	0.9989	1.4681	3
Q 0119–286	0.1170	0.0080	0.1058	3
Q 0119–013	0.0540	0.0080	0.0435	3
Q 0123+257	2.37	1.644	2.338	16
PC 0131+0120	3.792	3.116	3.744	1
Q 0132–1947	3.130	1.714	3.089	21
Q 0134+329	0.3670	0.0080	0.1050	3
BRI B0135–4239	3.97	2.575	3.920	1
Q 0136+010	2.35	1.749	2.317	16
Q 0136+1737	2.73	1.632	2.679	21
Q 0143–0135	3.141	1.673	3.097	21
GB 0148+2502	3.10	1.825	3.059	2
Q 0148–0946	2.850	1.797	2.810	21
BRI B0151–0025	4.194	2.74	4.142	22
Q 0153+0430	2.993	1.673	2.951	21
Q 0157+001	0.1631	0.0080	0.1515	3
Q 0159+036	2.47	1.644	2.436	16
Q 0205+024	0.1564	0.0080	0.1448	3
Q 0207–0019	2.853	1.756	2.817	21
Q 0215+015	1.7150	0.9996	1.5500	3
Q 0219+428	0.4440	0.0080	0.4296	3
GB 0229+1309	2.07	1.767	2.039	2
Q 0232–042	1.4360	0.0080	0.6320	3
–	–	0.8733	1.4116	3
Q 0237–233	2.2230	1.1593	1.5402	3

TABLE 6—*Continued*

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
Q 0239–1527	2.786	1.928	2.744	21
BRI B0241–0146	4.053	2.86	4.002	22
BR B0245–0608	4.238	2.96	4.186	22
PSS J0248+1802	4.43	2.810	4.376	1
Q 0249–1826	3.210	1.871	3.163	21
Q 0249–2212	3.21	2.044	3.160	5,1
Q 0252+0136	2.47	1.634	2.430	4
Q 0254+0000	2.25	1.634	2.215	4
Q 0256–0000	3.377	2.241	3.330	21
Q 0256–0031	2.00	1.634	1.965	4
Q 0258+0210	2.52	1.634	2.489	4
Q 0301–0035	3.226	2.060	3.181	21
Q 0302–0019	3.290	1.739	3.243	21
Q 0302–223	1.4000	1.0077	1.3760	3
Q 0305+0127	2.15	1.634	2.118	4
Q 0307–0058	2.11	1.634	2.075	4
Q 0308+0129	2.34	1.739	2.302	4
Q 0308+1902	2.839	1.673	2.797	21
Q 0308–1920	2.756	1.673	2.714	21
Q 0312–770	0.2230	0.0080	0.2108	3
Q 0316–2023	2.869	1.747	2.826	21
Q 0323+022	0.1470	0.0080	0.1295	3
Q 0329–2534	2.689	1.661	2.662	21
Q 0334–2029	3.132	2.057	3.089	21
PC 0345+0130	3.638	2.699	3.592	1
BR B0351–1034	4.351	3.09	4.297	22
Q 0351–3904	3.01	1.632	2.970	21
Q 0352–2732	2.823	1.673	2.781	21
BR B0401–1711	4.236	2.82	4.184	22
Q 0405–123	0.5740	0.0080	0.5583	3
Q 0414–060	0.7810	0.0080	0.7632	3
Q 0420+007	2.918	1.673	2.879	21
Q 0420–3851	3.1230	2.094	3.082	21
Q 0428–1342	3.244	1.965	3.200	21
Q 0454–220	0.5340	0.1199	0.5187	3
Q 0454+039	1.3450	0.9672	1.3216	3
Q 0457+024	2.38	1.645	2.346	16
MG 0504+0303	2.46	1.803	2.425	2
Q 0521–365	0.0566	0.0080	0.0460	3
Q 0537–441	0.8940	0.5139	0.6300	3
Q 0548–322	0.0690	0.0080	0.0583	3
Q 0552+398	2.36	1.644	2.325	16
Q 0558–504	0.1370	0.0080	0.1256	3
Q 0624+691	0.3700	0.0080	0.3563	3
Q 0636+6801	3.178	2.019	3.132	21

TABLE 6—*Continued*

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
Q 0637–752	0.6560	0.0080	0.6251	3
Q 0642+4454	3.408	2.192	3.362	21
Q 0702+646	0.0795	0.0080	0.0687	3
Q 0731+6519	3.038	2.019	2.993	21
Q 0735+178	0.4240	0.0765	0.4098	3
Q 0736+017	0.1910	0.0080	0.1791	3
Q 0742+318	0.4620	0.0080	0.4474	3
Q 0743–673	1.5130	1.0302	1.4879	3
GB 0749+4239	3.59	2.185	3.544	2
PC 0751+5623	4.281	3.526	4.228	1
Q 0754+100	0.6700	0.0080	0.6257	3
Q 0754+394	0.0958	0.0080	0.0848	3
Q 0804+761	0.1000	0.0080	0.0890	3
Q 0805+0441	2.880	1.838	2.834	21
Q 0812+332	2.42	1.677	2.385	16
Q 0819–032	2.35	1.704	2.319	16
Q 0820+296	2.37	1.644	2.333	16
MG 0830+1009	3.75	2.040	3.703	2
Q 0830+1133	2.979	1.797	2.936	21
Q 0831+1238	2.748	1.961	2.706	21
Q 0837–120	0.1980	0.0080	0.1860	3
Q 0844+349	0.0640	0.0080	0.0534	3
Q 0846+152	2.64	1.831	2.599	16
MG 0848+1533	2.01	1.735	1.980	2
Q 0849+080	0.0620	0.0080	0.0514	3
Q 0851+202	0.3060	0.0080	0.2929	3
Q 0855+182	2.62	1.682	2.580	16
Q 0903+155	2.68	1.659	2.645	16
MG 0906+0406	3.20	1.811	3.158	2
Q 0906+484	0.1180	0.0080	0.1068	3
Q 0910+403	0.9360	0.0080	0.9166	3
Q 0914–621	0.0573	0.0080	0.0467	3
Q 0916+555	0.1235	0.0080	0.1123	3
Q 0932+3646	2.84	1.634	2.814	21
Q 0933+733	2.53	1.651	2.493	16
Q 0938+1159	3.19	1.634	3.149	21
Q 0941+2608	2.913	1.731	2.867	21
Q 0953+414	0.2390	0.0080	0.2266	3
Q 0955+326	0.5330	0.0080	0.5177	3
Q 0956+1217	3.306	2.159	3.263	21
Q 0957+561	1.4050	0.8179	1.3810	3
Q 0958+551	1.7324	1.1762	1.4513	3
Q 1001+291	0.3290	0.0080	0.3157	3
Q 1004+130	0.2410	0.0080	0.2286	3
Q 1004+1411	2.707	1.786	2.672	21

TABLE 6—*Continued*

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
Q 1007+417	0.6110	0.0080	0.5949	3
Q 1009−0252	2.75	1.651	2.708	4
Q 1011−282	0.6110	0.0080	0.1310	3
Q 1011−0144	2.24	1.669	2.204	4
Q 1011+250	1.6310	0.9718	1.5500	3
Q 1012+008	0.1850	0.0080	0.1732	3
Q 1012−0206	2.14	1.634	2.104	4
GB 1013+2052	3.11	1.945	3.069	2
Q 1014+0023	2.29	1.634	2.591	4
Q 1016−0039	2.18	1.649	2.144	4
Q 1017+1055	3.158	2.114	3.127	21
Q 1017+280	1.9280	0.9971	1.4678	3
Q 1018−0005	2.60	1.789	2.560	4
Q 1020+0028	1.90	1.680	1.872	4
Q 1021−0037	2.547	1.887	2.513	21
Q 1024+0030	2.17	1.717	2.135	4
Q 1025−0030	2.87	1.885	2.833	4
Q 1028+313	0.1770	0.0080	0.1652	3
Q 1029−140	0.0860	0.0080	0.0751	3
Q 1033+1342	3.07	1.800	3.048	21
BR B1033−0327	4.509	2.91	4.454	22,23
Q 1038+528	2.30	1.677	2.262	16
GB 1041+3014	2.99	1.735	2.950	2
Q 1047+550	2.1650	1.3299	1.5159	3
BRI B1050−0000	4.286	2.83	4.233	22
Q 1100+772	0.3110	0.0080	0.2979	3
Q 1100−264	2.1450	1.1551	1.5500	3
MG 1101+0248	2.51	1.736	2.475	2
Q 1103−006	0.4260	0.0080	0.4117	3
BRI B1110+0106	3.918	2.58	3.869	22
Q 1115+080	1.7180	0.4066	0.6330	3
−	−	0.9595	1.5500	3
Q 1116+215	0.1770	0.0080	0.1652	3
Q 1123+264	2.35	1.645	2.317	16
Q 1124+5706	2.890	1.762	2.851	21
Q 1127+078	2.66	1.644	2.621	16
Q 1128+105	2.65	2.040	2.610	16
Q 1131−0043	2.16	1.653	2.128	4
Q 1132−0054	2.76	1.717	2.718	4
Q 1135−0255	2.41	1.739	2.373	4
Q 1136−135	0.5570	0.0080	0.5414	3
Q 1136+122	2.90	1.781	2.862	16
Q 1137+660	0.6460	0.0080	0.6295	3
Q 1138−0107	2.76	1.953	2.718	4
Q 1139−0139	1.93	1.634	1.884	4

TABLE 6—*Continued*

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
Q 1139−0037	1.91	1.634	1.896	4
Q 1142+0138	2.42	1.791	2.390	4
Q 1142+1015	3.152	2.127	3.109	21
Q 1143+0142	2.28	1.634	2.248	4
Q 1143+099	2.60	1.676	2.567	16
Q 1144+115	2.51	1.682	2.471	16
Q 1144+0140	2.59	1.667	2.551	4
Q 1145−0039	1.94	1.634	1.912	4
Q 1145+0121	2.08	1.721	2.045	4
Q 1146+0207	2.06	1.634	2.025	4
Q 1147+084	2.61	1.854	2.577	16
GB 1147+4348	3.02	2.035	2.980	2
Q 1148−0007	1.977	1.634	1.947	4
Q 1148+0055	1.89	1.667	1.858	4
Q 1148+549	0.9690	0.0080	0.9493	3
Q 1151+117	0.1760	0.0080	0.1642	3
Q 1156+295	0.7290	0.0080	0.7117	3
Q 1159+0039	2.586	1.671	2.550	21
Q 1202+281	0.1650	0.0080	0.1534	3
Q 1205−3014	3.036	2.045	2.996	21
Q 1206+1155	3.106	2.039	3.073	21
Q 1206+1500	2.60	1.793	2.568	4
Q 1206+1727	2.36	1.634	2.321	4
Q 1206+459	1.1580	0.4231	0.6300	3
—	—	0.8426	1.1364	3
Q 1209+1046	2.20	1.634	2.163	4
Q 1209+1524	3.06	1.634	3.021	4
Q 1211+143	0.0850	0.0080	0.0742	3
Q 1212+1551	1.95	1.665	1.918	4
Q 1212+1045	1.95	1.634	1.922	4
Q 1212+0854	2.35	1.634	2.319	4
Q 1213+1015	2.52	1.634	2.482	4
Q 1213+0922	2.72	1.675	2.681	4
Q 1215+1244	2.08	1.634	2.048	4
Q 1215+1202	2.83	1.634	2.788	4
Q 1215+303	0.2370	0.0080	0.2246	3
Q 1216+069	0.3340	0.0080	0.3207	3
Q 1216+1517	1.83	1.723	1.802	4
Q 1216+1754	1.81	1.634	1.781	4
Q 1216+1656	2.83	1.659	2.791	4
Q 1216+0947	2.31	1.645	2.279	4
Q 1217+023	0.2400	0.0080	0.2276	3
Q 1218+304	0.1300	0.0080	0.1187	3
Q 1219+755	0.0700	0.0080	0.0593	3
Q 1219+1140	2.18	1.634	2.147	4
Q 1222+228	2.0400	0.4647	0.6316	3

TABLE 6—*Continued*

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
Q 1222+1053	2.30	1.641	2.263	4
Q 1223+1059	2.32	1.643	2.288	4
Q 1223+1723	2.42	1.659	2.386	4
Q 1224+1244	2.14	1.634	2.110	4
Q 1225+1512	2.01	1.797	1.977	4
Q 1225+1610	2.23	1.663	2.200	4
Q 1225+317	2.2190	1.1263	1.5500	3
Q 1226+1035	2.32	1.634	2.287	4
Q 1226+1115	1.98	1.634	1.950	4
Q 1226+1639	2.25	1.634	2.216	4
Q 1226+023	0.1580	0.0080	0.1464	3
Q 1227+1215	2.17	1.624	2.138	4
Q 1228+1808	2.64	1.780	2.607	4
Q 1228+077	2.39	1.691	2.354	16
Q 1229+1414	2.90	1.764	2.862	4
Q 1229+1531	2.27	1.634	2.237	4
Q 1229−021	1.0380	0.4738	0.6320	3
Q 1229+204	0.0640	0.0080	0.0534	3
Q 1230+1042	2.43	1.634	2.396	4
Q 1230+1318	2.29	1.634	2.257	4
Q 1230+1627B	2.70	1.634	2.663	4
Q 1230+0941	1.84	1.641	1.812	4
Q 1232−0051	2.78	1.782	2.745	4
Q 1232+1139	2.87	1.848	2.831	4
Q 1234+0122	2.03	1.634	1.996	4
Q 1235+1807A	2.41	1.782	2.371	4
Q 1236−0043	1.84	1.690	1.815	4
Q 1236−0207	2.25	1.729	2.213	4
Q 1237+1515	2.04	1.634	2.009	4
Q 1237+0107	1.81	1.733	1.780	4
Q 1237+1508	2.07	1.634	2.035	4
Q 1237+1212	2.31	1.634	2.281	4
Q 1239+1435	1.93	1.634	1.900	4
Q 1239+0249	2.22	1.719	2.184	4
Q 1240+1504	1.85	1.634	1.823	4
Q 1241+176	1.2730	0.4066	0.6320	3
—	—	0.7657	1.2503	3
Q 1242+0213	1.99	1.634	1.958	4
Q 1242+0006	2.08	1.634	2.045	4
Q 1242+1732	1.83	1.696	1.805	4
Q 1242+1737	1.86	1.634	1.828	4
Q 1244+1129	3.16	2.101	3.118	4
Q 1244+1642	2.87	1.848	2.826	4
Q 1246−0059	2.45	1.669	2.415	4
Q 1246+0032	2.31	1.651	2.273	4
Q 1247+267	2.0380	0.9211	1.5500	3

TABLE 6—*Continued*

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
Q 1248+401	1.0300	0.3984	0.6028	3
—	—	0.8919	1.0097	3
Q 1253−055	0.5380	0.0080	0.5226	3
Q 1259+593	0.4720	0.0080	0.4573	3
Q 1302−102	0.2860	0.0080	0.2731	3
Q 1307+085	0.1550	0.0080	0.1435	3
Q 1308+326	0.9960	0.4670	0.6310	3
Q 1308−0214	2.85	1.892	2.811	4
Q 1308−0104	2.59	1.634	2.549	4
Q 1309+355	0.1840	0.0080	0.1722	3
Q 1312+043	2.35	1.813	2.319	16
Q 1313+0107	2.39	1.647	2.359	4
PSS J1317+3531	4.365	2.978	4.311	1
Q 1317+277	1.0220	0.2503	1.0018	3
Q 1318+290B	0.5490	0.3757	0.5335	3
Q 1318−0150	2.01	1.651	1.980	4
Q 1318−113	2.3080	1.896	2.273	16
Q 1320+0048	1.96	1.655	1.925	4
Q 1323−0248	2.12	1.661	2.090	4
Q 1324−0212	1.89	1.634	1.857	4
Q 1327−206	1.1690	1.1243	1.1473	3
Q 1328+0223	2.15	1.937	2.122	4
BRI B1328−0433	4.217	2.24	4.165	22
Q 1329+0231	2.43	1.663	2.400	4
Q 1329+0018	2.35	1.661	2.318	4
Q 1331+170	2.0840	1.2621	1.5500	3
Q 1333+176	0.5540	0.3902	0.5385	3
Q 1334+246	0.1070	0.0080	0.0959	3
Q 1334−0033	2.78	1.634	2.745	4
Q 1334+0212	2.38	1.634	2.350	4
BRI B1335−0417	4.396	3.08	4.342	22
Q 1336+0210	1.96	1.634	1.932	4
GB 1338+3809	3.10	1.737	3.059	2
Q 1338+101	2.45	1.724	2.412	16
Q 1338+416	1.2190	0.4066	0.6324	3
—	—	0.8684	1.1968	3
Q 1340+0959	2.942	1.894	2.897	21
Q 1344+0137	1.92	1.634	1.886	4
Q 1345−0137	1.93	1.634	1.900	4
Q 1345−0120	2.95	1.926	2.906	4
Q 1346+0121A	1.93	1.634	1.901	4
Q 1346−036	2.36	1.653	2.327	16
Q 1351+640	0.0880	0.0080	0.0771	3
Q 1352+183	0.1520	0.0080	0.1405	3
Q 1352+108	3.18	1.928	3.137	16
Q 1353+186	0.0505	0.0080	0.0400	3

TABLE 6—*Continued*

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
Q 1354+195	0.7200	0.3593	0.6330	3
Q 1355−416	0.3130	0.0080	0.2999	3
Q 1356+581	1.3710	0.5218	0.6310	3
Q 1358+115	2.59	1.677	2.550	16
Q 1358+3908	3.3	2.221	3.237	21
Q 1400+0935	2.980	2.022	2.930	21
Q 1402−012	2.52	1.789	2.479	16
Q 1402+044	3.20	2.340	3.160	16
Q 1406+123	2.94	2.018	2.903	16
Q 1407+265	0.9440	0.0080	0.9246	3
Q 1410+096	3.21	2.099	3.169	16
Q 1411+442	0.0900	0.0080	0.0791	3
GB 1413+3720	2.36	1.735	2.326	2
Q 1415+451	0.1140	0.0080	0.1029	3
Q 1416−129	0.1290	0.0080	0.1177	3
Q 1418+546	0.1520	0.0080	0.1405	3
Q 1419+480	0.0720	0.0080	0.0613	3
Q 1421+330	1.9040	1.0311	1.5500	3
Q 1425+267	0.3620	0.2409	0.3484	3
Q 1426+015	0.0860	0.0080	0.0751	3
Q 1428+0202	2.11	1.634	2.075	4
Q 1429−0053	2.08	1.719	2.047	4
Q 1429+118	3.00	1.958	2.963	16
PSS J1430+2828	4.306	2.777	4.253	1
Q 1433+0223	2.14	1.634	2.111	4
Q 1433−0025	2.04	1.634	2.012	4
PSS J1435+3057	4.297	2.905	4.244	1
GB 1436+4431	2.10	1.769	2.069	2
Q 1439+0047	1.86	1.649	1.828	4
Q 1440−0024	1.81	1.634	1.786	4
Q 1440+356	0.0781	0.0080	0.0673	3
Q 1444+407	0.2670	0.0080	0.2543	3
Q 1444+0126	2.21	1.717	2.174	4
Q 1444−0112	2.15	1.651	2.121	4
Q 1451−375	0.3140	0.0080	0.3009	3
Q 1455+123	3.08	1.830	3.033	16
MG 1500+0431	3.67	2.606	3.623	1
Q 1503+118	2.78	1.957	2.740	16
GB 1508+5714	4.283	2.73	4.230	22
Q 1512+370	0.3710	0.0080	0.3573	3
MG 1519+1806	3.06	1.955	3.019	2
GB 1520+4347	2.18	1.775	2.148	2
Q 1522+101	1.3210	0.0080	0.6310	3
—	—	0.8803	1.2978	3
Q 1525+227	0.2530	0.0080	0.2405	3
GB 1526+6701	3.02	1.955	2.980	2

TABLE 6—*Continued*

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
Q 1526+285	0.4500	0.0080	0.2428	3
Q 1538+477	0.7700	0.3326	0.6326	3
Q 1545+210	0.2640	0.0080	0.2514	3
Q 1548+0917	2.749	1.874	2.707	21
PC 1548+4637	3.544	2.607	3.499	1
Q 1553+113	0.3600	0.0080	0.3464	3
Q 1556+273	0.0899	0.0080	0.0790	3
MG 1557+0313	3.891	2.66	3.842	22
MG 1559+1405	2.24	1.737	3.059	2
Q 1600+0729	4.38	3.062	4.326	1
Q 1607+1819	3.123	1.814	3.0918	21
Q 1612+261	0.1310	0.0080	0.1197	3
Q 1613+658	0.1290	0.0080	0.1177	3
Q 1623+268A	2.47	1.644	2.433	16
Q 1623+268B	2.54	1.644	2.502	16
Q 1630+377	1.4710	0.0080	0.6320	3
—	—	0.8641	1.4463	3
Q 1631+3722	2.940	1.785	2.906	21
Q 1634+706	1.3340	0.5547	1.3107	3
Q 1641+399	0.5950	0.0080	0.5791	3
PC 1640+4628	3.700	2.604	3.653	1
Q 1704+608	0.3710	0.0080	0.3573	3
Q 1705+0152	2.576	1.669	2.537	21
Q 1715+535	1.9290	1.1009	1.5500	3
Q 1718+481	1.0840	0.0080	1.0632	3
Q 1721+343	0.2060	0.0080	0.1939	3
Q 1726+3425	2.429	1.669	2.393	21
Q 1727+502	0.0550	0.0080	0.0445	3
Q 1738+3502	3.240	2.093	3.197	21
GB 1745+6227	3.901	2.47	3.852	22
Q 1803+676	0.1360	0.0080	0.1246	3
Q 1807+698	0.0512	0.0080	0.0407	3
Q 1821+643	0.2970	0.0080	0.2840	3
Q 1831+731	0.1230	0.0080	0.1118	3
Q 1833+326	0.0590	0.0080	0.0484	3
Q 1836+5108	2.827	1.920	2.789	21
Q 1839–785	0.0743	0.0080	0.0636	3
Q 1845+797	0.0556	0.0080	0.0450	3
Q 1912–550	0.4020	0.1769	0.2041	3
Q 1928+738	0.3020	0.0080	0.2890	3
PKS 1937–101	3.787	2.442	3.739	21
Q 2000–3300	3.783	2.521	3.729	21
Q 2005–489	0.0710	0.0080	0.0603	3
Q 2038–0116	2.783	1.887	2.745	21
Q 2045–377	1.8000	1.0040	1.5500	3
Q 2048+3116	3.198	1.830	3.143	21

TABLE 6—*Continued*

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
Q 2050–359	3.49	2.605	3.445	1
Q 2113–4345	2.05	1.664	2.023	4
Q 2113–4534	2.54	1.969	2.506	4
Q 2114–4346	2.04	1.606	2.011	4
Q 2115–4434	2.16	1.755	2.128	4
Q 2117–4703	2.26	1.849	2.223	4
Q 2122–4231	2.27	1.550	2.233	4
Q 2126–1551	3.2660	2.011	3.218	21
Q 2126–4618	1.89	1.715	1.859	4
Q 2127–4528	2.71	2.018	2.676	4
Q 2128–123	0.5010	0.0940	0.4860	3
Q 2130+099	0.0610	0.0080	0.0504	3
Q 2131–4257	2.10	1.590	2.065	4
Q 2134–4239	1.80	1.590	1.776	4
Q 2134–147	0.2000	0.0080	0.1880	3
Q 2135–4632	2.21	1.879	2.182	4
Q 2136+141	2.43	1.784	2.390	16
Q 2139–4434	3.23	2.373	3.188	4
Q 2141+175	0.2130	0.0080	0.2009	3
Q 2145+067	0.9900	0.9426	0.9701	3
MG 2152+1420	2.56	1.800	2.524	2
Q 2153–2056	1.85	1.634	1.821	4
Q 2155–304	0.1170	0.0080	0.1058	3
Q 2159–2058	2.12	1.634	2.089	4
Q 2201+315	0.2970	0.0080	0.2840	3
Q 2203–2145	2.27	1.692	2.240	4
Q 2203–1833	2.73	1.849	2.691	4
Q 2205–2014	2.64	1.652	2.599	4
MG 2206+1753	3.14	1.769	3.099	2
Q 2209–1842	2.09	1.634	2.061	4
Q 2209+184	0.0700	0.0080	0.0593	3
Q 2211–1915	1.95	1.634	1.923	4
BR B2212–1626	3.990	2.69	3.940	22
Q 2214+139	0.0658	0.0080	0.0551	3
MG 2222+0511	2.32	1.800	2.287	2
GB 2223+2024	3.56	2.101	3.514	2
Q 2231+0125	1.90	1.634	1.871	4
Q 2231–0212	1.90	1.634	1.871	4
Q 2233+1341	3.209	2.216	3.167	21
Q 2233+1310	3.298	2.134	3.252	21
Q 2241+0014	2.14	1.657	2.099	4
Q 2243+0141	2.30	1.663	2.267	4
Q 2244–0234	1.97	1.787	1.940	4
Q 2244–0105	2.04	1.634	2.010	4
Q 2246–0006	2.05	1.651	2.019	4
BR B2248–1242	4.161	2.94	4.109	22

TABLE 6—*Continued*

QSO	z_{em}	z_{min}	z_{max}	Survey Reference
MG 2251+2429	2.33	2.019	2.297	2
Q 2251–178	0.0680	0.0080	0.0573	3
Q 2251+113	0.3230	0.1310	0.3098	3
MG 2254+0227	2.09	2.767	2.059	2
Q 2256+017	2.67	1.786	2.629	16
Q 2302+029	1.0440	0.3942	0.6290	3
–	–	0.8060	1.0236	3
Q 2308+098	0.4320	0.0080	0.4177	3
Q 2311–0341	3.048	1.714	3.001	21
MG 2320+0755	2.09	1.780	2.059	2
Q 2326–477	1.2990	0.9164	1.2760	3
PC 2331 + 0216	4.093	3.115	4.042	1
Q 2334+1041	2.243	1.634	2.211	21
Q 2344+092	0.6720	0.0080	0.6288	3
Q 2351+1042	2.379	1.632	2.345	21
Q 2351+0120	2.07	1.634	2.039	4
Q 2351–1154	2.67	1.632	2.633	21
Q 2352+0205	2.19	1.634	2.158	4
Q 2354–0134	2.21	1.665	2.178	4
Q 2356+0139	2.07	1.661	2.039	4
Q 2356+0237	2.50	1.634	2.465	4
Q 2359+0653	3.238	1.632	3.203	21
Q 2359+0023	2.897	1.714	2.857	21

REFERENCES.— (1) this paper; (2) Storrie-Lombardi & Hook 2000; (3) Lanzetta, Wolfe, & Turnshek 1995; (4) WLF95; (5) Sargent, Steidel, & Boksenberg 1989; (6) Turnshek *et al.* 1989; (7) Wolfe *et al.* 1993; (8) Lu *et al.* 1993; (9) Lu & Wolfe; (10) Virgilio *et al.* 1995; (11) Pettini *et al.* 1994; (12) Francis & Hewett 1993; (13) Savaglio *et al.* 1994; (14) Sargent, Boksenberg, & Steidel 1988; (15) Black, Chaffee, & Foltz 1987; (16) Wolfe *et al.* 1986; (17) Wolfe *et al.* 1994; (18) Rauch *et al.* 1990; (19) Williger *et al.* 1989; (20) Meyer, Lanzetta & Wolfe 1995; (21) Lanzetta *et al.* 1991; (22) SMH96; (23) SIM96

TABLE 7
REDSHIFT PATH SURVEYED

Redshift Range	Survey	Δz	ΔX		# of QSOs
			$q_0=0$	$q_0=0.5$	
3.00 - 5.00	This Paper	73.8	332.5	156.5	128
	APM	43.5	195.8	92.2	82
	LBQS	13.4	56.3	27.5	61
0.01 - 5.00	This Paper	418.4	1364.2	747.6	645
	APM	239.5	764.4	421.3	366
	LBQS	323.8	966.8	554.1	571

REFERENCES.—APM = SMIH96; LBQS = WLFC95

TABLE 8
FIT TO HI COLUMN DENSITY DISTRIBUTION

$f(N,z)=(f_*/N_*) (N/N_*)^{-\beta} e^{-N/N_*}$				
Redshift Range	f_*	$\log N_*$	β	
Including Lyman-limit System Data				
1.5 - 4.0	0.051	$21.22^{+0.10}_{-0.16}$	$1.24^{+0.02}_{-0.08}$	
4.0 - 4.7	0.160	$21.22^{+0.52}_{-0.33}$	$1.08^{+0.14}_{-0.22}$	
Not Including Lyman-limit System Data				
1.5 - 4.0	0.03	21.39 ± 0.30	1.52 ± 0.25	
4.0 - 4.7	0.34	20.92 ± 0.50	0.32 ± 0.60	

TABLE 9
DATA FOR FIGURE 14

Redshift Range	DLA $\langle z \rangle$	Δz	Number of DLAs QSOs		$\Omega = 1, \Lambda = 0$			$\Omega = 0.3, \Lambda = 0$			$\Omega = 0.3, \Lambda = 0.7$		
					ΔX	Ω_g	σ	ΔX	Ω_g	σ	ΔX	Ω_g	σ
.008 – 1.50	0.65	47.8	4	178	58.5	0.54	0.31	67.1	0.47	0.27	80.7	0.39	0.22
1.50 – 2.00	1.84	96.4	19	372	162.1	1.27	0.36	219.1	0.94	0.27	281.8	0.73	0.21
2.00 – 2.50	2.26	124.2	25	349	223.0	1.67	0.56	310.3	1.20	0.40	393.7	0.94	0.32
2.50 – 3.00	2.69	76.9	15	234	148.4	1.54	0.51	212.6	1.08	0.36	265.1	0.86	0.29
3.00 – 3.50	3.21	40.9	10	128	83.9	2.10	0.80	123.0	1.44	0.54	151.0	1.17	0.44
3.50 – 4.70	3.98	33.8	12	60	74.5	1.23	0.43	111.8	0.82	0.28	134.6	0.68	0.24

NOTE.—When totalled, the number of QSOs in column 5 is more than the total for this data set (646). This is because the same QSO can contribute redshift path to more than one redshift bin.